


SOLVED EXAMPLES

JEE Main/Boards

Example 1: A 10.0 litre vessel contains 2 moles of B at 100 K. How many moles of C should be added to drive the following backward reaction practically to completion?



Sol: Using the given K_p , calculate the partial pressure of gas C and use it to get the no. of moles. Using the reaction, moles of C can be found.

$$K_p = [p_c]^2 = 16; P_c = 4 \text{ atm}$$

Using ideal gas equation, for C,

$$pV = nRT$$

$$4 \times 10 = n_c \times 0.0821 \times 100$$

$$n_c = \frac{0.0821 \times 100}{4 \times 10}$$




$$n_c = 4.872 \text{ moles}$$

By looking at the given equation, stoichiometry for B and C should be in the ration 1:2

Thus 2 moles of B shall combine with 4 moles of C to give back A.

In order to drive the backward reaction almost to completion total number of moles required will be = 4.872 + 4 = 8.872

Example 2: (a) For which of the following reactions, K_p is equal to K_c ?



Sol: Use equation $K_p = K_c(RT)^{\Delta n}$ to solve the above.

In order to calculate K_p , we have to use the formula, $K_p = K_c(RT)^{\Delta n}$

For reaction $H_2 + I_2 \rightleftharpoons 2HI$ only $\Delta n = 2 - 2 = 0$

For this reaction $(RT)^{\Delta n} = 0$

Thus $K_p = K_c$



Example 3: For which of the following cases does the reaction go farthest to completion:

$$K = 1, K = 10^{10}, K = 10^{-10}?$$

Sol: More the value of the ratio of [product]/[reactant], farther the reaction will go.

The reaction having $K=10^{10}$ will go farthest to completion because the ratio [product]/[reactant] is maximum in this case.

Example 4: Both metal Mg and Fe can reduce the metal copper from a solution having copper ions (Cu^{2+}), according to the equilibria:



Which metal will remove cupric ions from the solution to a greater extent?

Sol:


- As we know greater the value of K forward reaction is more feasible.
- Since $K_1 > K_2$, the product in the first reaction is much more favoured than in the second one.
- Mg thus removes more Cu^{2+} from solution than does Fe.

(K_1 and K_2 include the constant concentration of the solid species.)

Example 5: Two moles of PCl_5 were heated to $327^\circ C$ in a closed two litre vessel and when equilibrium was achieved, PCl_5 was found to be 40% dissociated into PCl_3 and Cl_2 . Calculate the equilibrium constants K_p and K_c for this reaction,

Sol: Amount of reacting species and the product can be calculated and then apply the 40% dissociation. Accordingly solve K_p and K_c by using $K_p = K_c(RT)^{\Delta n}$

Reaction for dissociation of PCl_5 :



Given: Amount of PCl_5 (initially) = 2 moles

Percentage dissociation at equilibrium = 40%

∴ No. of moles of PCl_5 dissociated at equilibrium

$$= \frac{40}{100} \times 2 = 0.8 \text{ mole}$$

∴ Amounts of PCl_5 , PCl_3 and Cl_2 at equilibrium will be

$$\text{PCl}_5 = 2 - 0.8 = 1.2 \text{ mole}$$

[From the reaction, 1 mole of PCl_5 on dissociation gives 1 mole of PCl_3 and 1 mole of Cl_2]

$$\therefore \text{PCl}_3 = 0.8 \text{ mole}$$

$$\therefore \text{Cl}_2 = 0.8 \text{ mole}$$

In order to calculate K_c we have to calculate molar concentration of reactant and product at equilibrium.

(Given: volume of vessel = 2 litre)

Thus,

$$[\text{PCl}_5] = \frac{1.2}{2} = 0.6 \text{ mol L}^{-1},$$

$$[\text{PCl}_3] = \frac{0.8}{2} = 0.4 \text{ mol L}^{-1}$$

$$\text{And } [\text{Cl}_2] = \frac{0.8}{2} = 0.4 \text{ mol L}^{-1}$$

Applying the law of chemical equilibrium to the dissociation equilibrium, we get

$$K_c = \frac{[\text{PCl}_3][\text{Cl}_2]}{[\text{PCl}_5]} = \frac{0.4 \text{ mol L}^{-1} \times 0.4 \text{ mol L}^{-1}}{0.6 \text{ mol L}^{-1}}$$

$$K_c = 0.267 \text{ mole L}^{-1}$$

As we have calculated the value of K_c we can now calculate value of K_p using the following equation.

$$K_p = K_c(RT)^{\Delta n}$$

$$\text{Now, } \Delta n = n_p = n_r = 2 - 1 = 1 \text{ mole}$$

$$\therefore K_p = K_c(RT)$$

$$T = 327 + 273 = 600 \text{ K}$$

$$R = 0.0821 \text{ L atm K}^{-1} \text{ mol}^{-1}$$


$$\therefore K_p = 0.267 \text{ mol L}^{-1} \times 0.0821 \text{ L atm K}^{-1} \text{ mol}^{-1} \times 600 \text{ K} = 13.15 \text{ atm}$$

Example 6: AB_2 dissociates as: $\text{AB}_2(\text{g}) \rightleftharpoons \text{AB}(\text{g}) + \text{B}(\text{g})$.

If the initial pressure is 500 mm of Hg and the total pressure at equilibrium is 700 mm of Hg, calculate K_p for the reaction.

Sol: Using partial pressure concepts in the given reaction, calculate the partial pressure and then apply K_p expression.

After dissociation, suppose the decreases in the pressure of AB_2 at equilibrium is p mm. Then



$$PT = 500 - p + p + p = 500 + p$$

$$\therefore 700 = 500 + p \therefore p = 200 \text{ MM}$$


$$\therefore P_{\text{AB}_2} = 500 - 200 = 300 \text{ mm}$$

$$\therefore K_p = \frac{P_{\text{AB}} \times P_{\text{B}}}{P_{\text{AB}_2}} = \frac{200 \times 200}{300} = 133.33 \text{ mm}$$

Example 7: The degree of dissociation of PCl_5 at a certain temperature and pressure is 0.2. Calculate the pressure at which it will be half (50%) dissociated at the same temperature.

Sol: Using the reaction and the given degree of dissociation, frame the partial pressure equations and solve by using K_p .

Suppose α is the degree of dissociation, then



∴ Total number of moles at equilibrium

$$= 1 - \alpha + \alpha + \alpha = 1 + \alpha$$

If P is the total pressure at equilibrium, then partial pressures will be

$$p_{\text{PCl}_3} = \frac{\alpha}{1 + \alpha} P, \quad p_{\text{Cl}_2} = \frac{\alpha}{1 + \alpha} P,$$

$$p_{\text{PCl}_5} = \frac{1 - \alpha}{1 + \alpha} P; \quad K_p = \frac{p_{\text{PCl}_3} \times p_{\text{Cl}_2}}{p_{\text{PCl}_5}}$$

Substituting the values in above equation,

$$= \frac{\left(\frac{\alpha}{1 + \alpha} P\right) \times \left(\frac{\alpha}{1 + \alpha} P\right)}{\left(\frac{1 - \alpha}{1 + \alpha} P\right)}$$


$$K_p = \frac{\alpha^2}{1 - \alpha^2} P$$

We are given that at $P = 1 \text{ atm}$, $\alpha = 0.2$.

$$\text{Hence, } K_p = \frac{(0.2)^2}{1 - (0.2)^2} \times 1 \therefore K_p = 0.042$$

At 50%, dissociation i. e. $\alpha = 0.5$,

Suppose total pressure is P' . Then




$$0.042 = \frac{(0.5)^2}{1 - (0.5)^2} \times p$$

$$P' = 0.1272 \text{ atm}$$

Example 8: A vessel at 1000 K contains carbon dioxide with a pressure of 0.4 atm. Some of the carbon dioxide is converted to carbon monoxide on addition of graphite. Calculate the value of K if the total pressure at equilibrium is 0.7 atm.

Sol: Frame the reaction. Using the total pressure and the partial pressure concept, solve for K . The chemical reaction for the above transformation is:



Suppose decrease in pressure of CO_2 after reaction = p atm

Final pressure = 0.7

Then increase in pressure due to $CO = 2p$

Final pressure = $(0.4 - p) + 2p = 0.4 + p = 0.7$ atm



$\therefore p = 0.3$ atm. Hence, we have

$$p_{CO_2} = 0.4 - 0.3 = 0.1 \text{ atm}$$

$$p_{CO} = 2 \times 0.3 = 0.6 \text{ atm}$$

$$K = \frac{p_{CO}^2}{p_{CO_2}} = \frac{(0.6)^2}{0.1} = 3.6$$

Example 9: Two solids A and C dissociate into gas products as follows;





At 25° C, the pressure over excess solid A only is 400 atm, and that over solid C only is 60 atm. Find the pressure over solid mixture.

Sol: Arrange the stepwise conversion and using the given values of K_p , frame for partial pressure and solve for solid mixture pressure.

- When two solids A and C are taken together in a closed container, both decompose to give gases B, D and E.
- As D is the common gas, the dissociation of both the solids A and C shall be suppressed.

- Suppose that the partial pressures of B and D due to dissociation of only A are p_1 atm each, and the partial pressures of E and D due to dissociation of only C are p_2 atm each.



$$K_{p_1} = p_B \cdot p_D = p_1(p_1 + p_2) = 400 \quad \dots (i)$$

$$\text{and } K_{p_2} = p_E \cdot p_D = p_2(p_1 + p_2) = 900 \quad \dots (ii)$$

Solving equation. (1) and (2),


$$p_1 = 11.097 \text{ and } p_2 = 24.96 \text{ atm.}$$

$$\therefore \text{pressure over solid mixture} = 2(p_1 + p_2)$$

$$= 2(11.097 + 24.96) = 72.114 \text{ atm.}$$


Example 10: The equilibrium constant of the reaction, $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$ at 426° C is 65.5, what will be the value of the equilibrium constant

- if the reaction is reversed, and
- if the given reaction is represented as



Sol: Using $1/K$ and $(K)^n$, solve the above.

(a) The reverse reaction of the given reaction is



(Remember: If the reaction is reversed equilibrium constant is given by $1/k$)

$$\therefore \text{Equilibrium constant} = \frac{1}{65.5} = 0.0152$$

(b) The reaction $3H_2 + 3I_2 \rightleftharpoons 6HI$ has been obtained by multiplying the reaction $H_2 + I_2 \rightleftharpoons 2HI$ by 3.

(Remember: if the reaction is multiplied by n ,

then equilibrium constant = $(K)^n$)


$$\text{Hence, } K = (65.5)^3 = 281011.$$

Example 11: A mixture of 1.62 mol of N_2 , 1.20 mol of H_2 and 9.24 mole of NH_3 is introduced into a 20 L reaction vessel at 500K. At this temperature, the equilibrium constant, K_c for the reaction, $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$ is 1.7×10^2 .

Predict the direction of reaction.


Sol: Calculate the concentration of each species in the reaction and then solve for Q_c .

For a general reaction $aA + bB \rightleftharpoons cC + dD$




$$Q = \frac{[\text{C}]^c [\text{D}]^d}{[\text{A}]^a [\text{B}]^b} = \frac{(0.462)^2}{(0.081)(0.08)^3} = 4.9 \times 10^3$$

For the given reaction, K_c



As $Q_c > K_c$, the net reaction will be in the backward direction.

Example 12: What is the effect of reducing the volume on the system described below?



Sol:

- This can be explained on the basis of Le Chatelier's principle.
- On reducing the volume, the pressure will increase.
- By Le Chatelier's principle equilibrium will shift to the side accompanied by decrease of pressure, i.e. decrease in the number of gaseous moles, i.e. backward direction.

Example 13: In the reaction $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$ at equilibrium, helium gas is injected into the vessel without disturbing the overall pressure of the system. What will be the effect on the equilibrium?

Sol: Application of Le-Chatelier's principle.

$$K_c = \frac{[\text{NH}_3]^2}{[\text{N}_2][\text{H}_2]^3}$$

- Since pressure is kept constant, volume will increase.
- Hence, molar concentration of NH_3 , N_2 and H_2 will decrease.
- As there are two concentration terms in numerator and four concentration terms in the denominator, to keep K_c constant, decrease in NH_3 should be more, i.e., equilibrium will shift in the backward direction.

Example 14: A liquid is in equilibrium with its vapour in a sealed container at fixed temperature. The volume of the container is suddenly increased.


- What is the initial effect of the change on vapour pressure?
- How do rates of evaporation and condensation change initially?
- What happens when equilibrium is restored finally and what will be the final vapour pressure?

Sol: (a) As volume is increased, the vapour pressure will decrease because the same amount of vapour are now distributed in larger space. Number of molecules striking the wall of container decreases and thus pressure decreases.

(b) As the vessel is sealed, the rate of evaporation remains constant at constant temperature. However, the rate of condensation will be low initially because there are fewer molecules per unit volume in the vapour phase and hence the number of collisions per unit time with the liquid surface decreases.

(c) When equilibrium is restored, Rate of evaporation = rate of condensation. The final vapour pressure = Initial pressure because vapour pressure of a liquid depends only on temperature and not volume.

Example 15: The following system is in equilibrium:



What will happen to the temperature of the system if some Cl_2 is added into it at constant volume?

Sol: Temperature will increase. Backward direction will be more feasible. Thus equilibrium will shift in the backward direction producing more heat.

Example 16: Explain why pure liquids and solids can be ignored while writing the equilibrium constant expression.

Sol: This can be explained by considering density of pure solid or pure liquid.


$$[\text{Pure liquid}] \text{ or } [\text{Pure solid}] = \frac{\text{No. of moles}}{\text{Volume in L}}$$

$$\begin{aligned} \text{No of moles} &= \text{Mass} / \text{Mol. Mass} = \frac{\text{Mass} / \text{mol.mass}}{\text{Volume}} \\ &= \frac{\text{Mass}}{\text{Volume}} \times \frac{1}{\text{Mol.mass}} = \frac{\text{Density}}{\text{Mol.mass}} \end{aligned}$$


- Density of a pure liquid or pure solid is constant at constant temperature and molecular mass is also constant, therefore, their molar concentrations are constant and are ignored into the equilibrium constant.

JEE Advanced/Boards

Example 1: One mole of nitrogen is mixed with 3 moles of hydrogen in 4-litre container. If 0.50 % of nitrogen is converted to ammonia by the following reaction,



Calculate the equilibrium constant (K_c) in concentration units. What will be the value of K_c for the following equilibrium?



Sol: Using the reaction, calculate equilibrium concentration and K_c . Apply the formula for the next reaction and solve the new K_c .

Initial moles	N_2	+	3H_2	\rightleftharpoons	$\text{NH}_3(\text{g})$
	1		3		0
Moles at equilibrium	(1-0.0050)		(3-0.0050)		(2×0.0050)
Molar concentration at equilibrium	$\frac{0.9950}{4}$		$\frac{2.9950}{4}$		$\frac{0.01}{4}$


$$K_c = \frac{[\text{NH}_3]^2}{[\text{N}_2][\text{H}_2]^3}$$

$$= \frac{\left(\frac{0.01}{4}\right)^2}{\left(\frac{0.9950}{4}\right)\left(\frac{2.9950}{4}\right)^3} = \frac{6.25 \times 10^{-6}}{(0.2487) \times (0.1044)}$$

$$= 6.0 \times 10^{-5} (\text{mol/L})^{-2}$$

For the other reaction $\frac{1}{2}\text{N}_2(\text{g}) + \frac{3}{2}\text{H}_2(\text{g}) \rightleftharpoons \text{NH}_3(\text{g})$

We can see that the above reaction has been obtained by multiplying the equation



Thus, the new equilibrium constant = $(6.0 \times 10^{-5})^{1/2}$

$$K_c = 7.7 \times 10^{-3} (\text{mole/litre})^{-1}$$

Example 2: Would 1% CO_2 in the air be sufficient to prevent any loss in weight when Ag_2CO_3 is dried at 120°C ? $\text{Ag}_2\text{CO}_3(\text{s})$ is dried at 120°C ? $\text{Ag}_2\text{CO}_3(\text{s}) \rightleftharpoons \text{Ag}_2\text{O}(\text{s}) + \text{CO}_2(\text{g})$; $K_p = 0.0095$ at 120°C . How low would the partial pressure of CO_2 have to be to promote this reaction at 120°C ?

Sol: $K_p = \frac{P_{\text{Ag}_2\text{O}} \times P_{\text{CO}_2}}{P_{\text{Ag}_2\text{CO}_3}} = P_{\text{CO}_2}$

$$\therefore P_{\text{CO}_2} = 0.0095 \text{ atm} = \text{constant at } 120^\circ\text{C}.$$

- Thus, if Ag_2CO_3 is taken in a closed container, a small amount of it would decompose to give CO_2 gas until the partial pressure of CO_2 reaches 0.0095 atm.
- As this is the equilibrium pressure of CO_2 , the decomposition would then stop.
- Now since partial pressure of CO_2 in air is 0.01 atm ($\because \text{CO}_2$ is 1% in air) which is much greater than 0.0095 atm, the equilibrium would practically shift to the left-hand side completely, or in other words, there would be no loss in weight of Ag_2CO_3 (by decomposition) if placed in air containing 1% CO_2 .
- Further, if the partial pressure of CO_2 in air is less than the equilibrium pressure of 0.0095 atm, the decomposition of Ag_2CO_3 would continue till the CO_2 pressure around Ag_2CO_3 becomes 0.0095 atm.

Example 3: In the preparation of quick lime from limestone, the reaction is



Experiments carried out between 850°C and 950°C led to a set of K_p values fitting an empirical equation

$$\log K_p = 7.282 - \frac{8500}{T}$$

where T is the absolute temperature. If the reaction is carried out in quiet air, what temperature would be predicted from this equation for complete decomposition of the limestone?

Sol: For the equilibrium $\text{CaCO}_3(\text{s}) \rightleftharpoons \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$

$$K_p = P_{\text{CO}_2}$$

the decomposition of CaCO_3 in quiet air, would continue till the pressure developed due to CO_2 equals 1 atm (atmospheric pressure).

\therefore when the decomposition is complete $K_p = 1 \text{ atm}$

Substituting K_p in the given empirical equation,

$$\log 1 = 7.282 - \frac{8500}{T} = 0.$$

$$T = 1167 \text{ K} = 894^\circ\text{C}$$

Example 4: 0.0785 g of selenium vapour occupying a volume of 105 mL of 600°C exerts a pressure of 160 mm. The selenium is in a state of equilibrium according to the reaction $\text{Se}_6(\text{g}) \rightleftharpoons 3\text{Se}_2(\text{g})$.

Calculate (a) degree of dissociation of selenium, (b) K_p , and (c) K_c ; ($\text{Se} = 79$)

Sol: Using $\left(\frac{p_1 V_1}{T_1} = \frac{p_2 V_2}{T_2}\right)$, find

1. Volume at NTP
2. No. of moles
3. Molar mass
4. Vapour density
5. Degree of dissociation
6. Using reaction, find moles at equilibrium and then solve for K_p and K_c from

$$K_p = K_c(RT)^{\Delta n}$$

$$(a) \text{ Volume at NTP} = \frac{160}{760} \times \frac{105}{873} \times 273 \text{ mL}$$

$$= 6.9216 \left(\frac{p_1 V_1}{T_1} = \frac{p_2 V_2}{T_2}\right)$$

$$\therefore \text{ Moles at NTP} = \frac{6.9216}{22400} \dots$$

Let the observed molecular weight of selenium be M.

$$\therefore \text{ No. of moles} = \frac{0.0785}{M}$$

$$\text{Hence, } \frac{0.0785}{M} = \frac{6.9216}{22400}$$

$$M = 254.04$$

$$\therefore \text{ Observed vapour density} = \frac{254.04}{2} = 127.02$$

$$\text{Theoretical vapour density} = \frac{79 \times 6}{2} = 237$$

(mol. wt. of $\text{Se}_6 = 79 \times 6$)

(i.e., when there is no diss.)

Since 1 molecule of selenium produces 3 molecules


$$\therefore n = 3$$

Substituting the above values in Eqn.(4), we get

$$\text{Degree of dissociation of } \text{Se}_6 = \frac{(237 - 127.02)}{(3 - 1) \times 127.02}$$

$$= \frac{190.98}{254.04} = 0.43$$

(b)



$$K_p = \frac{p_{\text{Se}_2}^3}{p_{\text{Se}_6}} = \frac{\left(\frac{3x}{1+2x} \times p\right)^3}{\left(\frac{1-x}{1+2x} \times p\right)} = \frac{27x^3}{(1+2x)^2(1-x)} \times p^2$$

Substituting the values of $x = 0.43$ and $p = \frac{160}{760}$ atm,

$$K_p = 0.8970$$

$$(c) K_p = K_c(RT)^{\Delta n} \dots \dots \text{(Eqn. 3)}$$

$$0.8970 = K_c (0.08821 \times 973)^2$$

$$(\Delta n = 3 - 1 = 2)$$

$$K_c = \frac{0.8970}{(0.821 \times 973)^2} = 1.40 \times 10^{-4}$$

Example 5: An equilibrium mixture,

$\text{CO}(\text{g}) + \text{H}_2\text{O}(\text{g}) \rightleftharpoons \text{CO}_2(\text{g}) + \text{H}_2(\text{g})$, present in a vessel of one litre capacity at 1000 K was found to contain 0.4 mole of CO, 0.3 mole of H_2O , 0.2 mole of CO_2 and 0.6 mole of H_2 . If it is desired to increase the concentration of CO to 0.6 mole by adding CO_2 into the vessel, how many moles of it must be added into equilibrium mixture at constant temperature in order to get this change?


Sol: Step-1 To calculate K_c of the reaction.

$$K_c = \frac{[\text{CO}(\text{g})][\text{H}_2(\text{g})]}{[\text{CO}(\text{g})][\text{H}_2\text{O}(\text{g})]} = \frac{0.2 \times 0.6}{0.4 \times 0.3} = 1$$

Step-2 To calculate extra CO_2 to be added:

Suppose X mole of extra CO_2 is added

Then writing the reverse reaction, we have



(V = 1 L) (Given)

$$K_c' = \frac{1}{K_c} = \frac{0.6 \times 0.5}{x \times 0.4} = 1 \quad \therefore x = \frac{0.6 \times 0.5}{0.4} = 0.75$$

Example 6: Ammonium carbamate decomposes as

$\text{NH}_2\text{COONH}_4(\text{s}) \rightleftharpoons 2\text{NH}_3(\text{g}) + \text{CO}_2(\text{g})$. In a closed vessel containing ammonium carbamate in equilibrium, NH_3 is added such that the partial pressure of NH_3 now

equals original total pressure. Calculate the ratio of total pressure now to the original pressure.

Sol: Let us assume that initial pressure of the mixture is P . Thus, pressure is due to NH_3 and CO_2 which are present in the ratio of 2 : 1. (from the reaction)

$$\text{Thus, } p_{\text{NH}_3} = \frac{2}{3}P, p_{\text{CO}_2} = \frac{1}{3}P$$

$$K_p = (p_{\text{NH}_3})^2 (p_{\text{CO}_2})^2 = \left(\frac{2}{3}P\right)^2 \left(\frac{1}{3}P\right) = \frac{4}{27}P^3$$

After adding NH_3 , partial pressure of NH_3 equals the original total pressure

$$p_{\text{NH}_3} = P \text{ (given)}$$


$$\therefore P^2 \times p_{\text{CO}_2}$$

$$= K_p = \frac{4}{27}P^3 \text{ or } p_{\text{CO}_2} = \frac{4}{27}P$$

$$\therefore \text{Total pressure now} = p_{\text{NH}_3} + p_{\text{CO}_2} = P + \frac{4}{27}P = \frac{31}{27}P = \frac{31}{27}$$

Example 7: An equilibrium mixture at 300 K contains N_2O_4 and NO_2 at 0.23 and 1.4 atm respectively. If the volume of the container is doubled, calculate the new equilibrium pressure of the two gases.

Sol: Step-1: Calculate K_p



$$\text{Equilibrium pressure } 0.23 \quad 1.4\text{atm}$$

$$K_p = \frac{p_{\text{NO}_2}^2}{p_{\text{N}_2\text{O}_4}} = \frac{(1.4\text{atm})^2}{0.23\text{atm}} = 8.52\text{atm}$$

Step-2: Calculation of new equilibrium pressures:

According to Boyle's law, volume increases pressure decreases.


Thus on doubling the volume, pressure will decrease to half.

As pressure decreases, equilibrium will shift to the side accompanied by increase in the number of mole, i.e. forward direction.

This means that pressure of N_2O_4 will decrease while that of NO_2 will increase

Let us assume that decrease in pressure of $\text{N}_2\text{O}_4 = p$.

Then



$$K_p = \frac{(0.70 + 2p)^2}{(0.115 - p)} = 8.52\text{atm} \text{ (Calculate above)}$$

$$0.49 + 4p^2 + 2.8p = 0.9798 - 8.52p$$

$$4p^2 + 11.32p - 0.4898 = 0$$

$$p = \frac{-11.32 \pm \sqrt{128.14 + 7.83}}{8} = 0.042\text{atm}$$

(minus value is neglected)

(For quadratic equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a})$$

\therefore New equilibrium pressures will be


$$p_{\text{N}_2\text{O}_4} = 0.115 - 0.042 = 0.072\text{atm},$$

$$p_{\text{NO}_2} = 0.70 + 2 \times 0.042 = 0.113\text{atm}$$

Example 8: At some temperature and under a pressure of 4 atm, PCl_5 is 20% dissociated. Calculate the pressure at which PCl_5 will be 40% dissociated, temperature remaining same.

Sol: Split into case 1 and case 2 where in case 1, 20% dissociation and in case 2, 40% dissociation is applied.

Case 1: When PCl_5 is 20% dissociated



$$\text{Total no. of moles} = 0.8 + 0.2 + 0.2 = 1.2\text{mole}$$


$$p_{\text{PCl}_5} = \frac{0.8}{1.2} \times 4\text{atm},$$

$$p_{\text{PCl}_3} = \frac{0.2}{1.2} \times 4\text{atm},$$

$$p_{\text{Cl}_2} = \frac{0.2}{1.2} \times 4\text{atm}$$

$$K_p = \frac{0.8 \times 0.8}{\frac{3.2}{1.2}} = 0.166$$

Case 2: When PCl_5 is 40% dissociated. Suppose total pressure = P atm. Then,



Total no. of moles = $0.6 + 0.4 + 0.4 = 1.4$ moles

$$p_{\text{PCl}_5} = \frac{0.6}{1.4} \times P \text{ atm}, p_{\text{PCl}_3} = \frac{0.4}{1.2} \times P \text{ atm}, p_{\text{Cl}_2} = \frac{0.4}{1.2} \times P \text{ atm}$$

$$K_p = \frac{\frac{0.4}{1.4} P \times \frac{0.4}{1.4} P}{\frac{0.6}{1.4} P} = \frac{0.4}{1.4} \times \frac{0.4}{0.6} P = 0.1904P$$

$$K_p = 0.166$$

Which gives $P = 0.87$ atm

$$\text{But } K_p = 0.166$$

$$\therefore P = 0.8 \text{ atm}$$

JEE Main/Boards

Exercise 1


Q.1 When 46 g of I_2 and Ig or H_2 are heated at equilibrium at 450°C , the equilibrium mixture contained 1.9g of I_2 . How many moles of I_2 and HI are present at equilibrium?

Q.2 A two litre flask contains 1.4 gm nitrogen and 1.0 gm hydrogen. The ratio of active mass of nitrogen and hydrogen would be?

Q.3 In the reaction: $\text{A} + \text{B} \rightleftharpoons \text{C} + \text{D}$ the initial concentration of A is double the initial concentration of B. At equilibrium the concentration of B was found to be one third of the concentration of C. The value of equilibrium constant is?

Q.4 The value of K_c for the reaction: $\text{A} + 3\text{B} \rightleftharpoons 2\text{C}$ at 400°C is 0.1768. Calculate the value of K_p ?

Q.5 Two moles of ammonia was introduced in an evacuated vessel of 1 litre capacity. At high temperature the gas undergoes particle dissociation according to the equation:




At equilibrium the concentration of ammonia was found be 1 mole. What is the value of 'K'?

Q.6 What will be the expressions of formation of PCl_5 for K_p and K_c ?

Q.7 4.0 gms of hydrogen react with 9.023×10^{23} molecules of chlorine to form HCl gas. The total pressure after the reaction was found to be 700 mm. What will be the partial pressure of HCl ?

Q.8 The equilibrium constant K for the reaction $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$ is 16. What would be value of equilibrium constant 'K' for the reaction:



Q.9 1.0 mole of PCl_3 (g) and 2.0 moles of Cl_2 (g) were placed in a 3 litre flask and heated to 400 K. When equilibrium was established, only 0.70 mole of PCl_3 (g) remained. What is the value of equilibrium constant for the reaction: $\text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g}) \rightleftharpoons \text{PCl}_5(\text{g})$ at 400 K?

Q.10 For the reaction: $\text{PCl}_5(\text{g}) \rightleftharpoons \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$

The moles of each component PCl_5 , PCl_3 and Cl_2 at equilibrium were found to be 2. If the total pressure is 3 atm. What will be the value of K_p ?


Q.11 For the reaction $\text{H}_2 + \text{I}_2 \rightleftharpoons 2\text{HI}$

The value of equilibrium constant is 9.0. Calculate the degree of dissociation of HI ?


Q.12 For the reaction $\text{N}_2 + \text{H}_2 \rightleftharpoons 2\text{NH}_3$, $\text{N}_2 : \text{H}_2$ where taken in the ratio of 1:3. Up to the points of equilibrium 50% each reactant has been reacted. If total pressure at equilibrium is P. Calculate the partial pressure of ammonia?

Q.13 In a reaction vessel of 2 litre capacity 3 mole N_2 reacts with 2 moles of O_2 to produce 1 mole of NO . What is the molar concentration of N_2 equilibrium?

Q.14 At 300 K, K_p for the reaction:



Calculate K_p and K_c for the reaction



Q.15 2.0 g mol of PCl_5 were heated in a 3 litre capacity vessel. At equilibrium 50% PCl_5 is dissociated. What will be the dissociation constant of the reaction?






Q.16 What should be the equilibrium constant K_2 for the reaction $2C \rightleftharpoons A + 3B$, if the equilibrium constant for the reaction $A + 3B \rightleftharpoons 2C$ is K_1 ?

Q.17 If the value of K_c for $P + Q \rightleftharpoons R + S$ is 10^{-2} and that of K_f is 10^{-1} the rate constant for the backward reaction will be.

Q.18 If the concentration of B is increased at fixed temperature, in the reaction, $A + 2B \rightleftharpoons C + 3D$ the equilibrium constant of backward reaction.

Q.19 In reaction $A + B \rightleftharpoons C + D$, if concentration of A is increased four times and concentration of B is halved, the rate of reaction will become.

Q.20 The value of for the reaction K_p^0 for the reaction $\frac{1}{2}H_2(g) + \frac{1}{2}I_2(g) \rightleftharpoons HI(g)$ is 8.32 at 873 K and 1 bare pressure. Calculate K_p^0 , K_c^0 and K_x for



Q.21 A mixture of hydrogen and iodine (molecular ratio is 2:1) is reacted to form HI. Calculate total moles at equilibrium?

Q.22 The vapour density for dissociation of PCl_5 at 250°C is 57.9. What will be the value of amount of dissociation, if molecular weight of PCl_5 is 208.5?

Q.23 In the reaction, $PCl_5 \rightleftharpoons PCl_3 + Cl_2$, a moles of PCl_5 are initially taken. If the amount x gets dissociated and total pressure is P, the value of $P_{PCl_3} \times P^{-1}$ will be

Q.24 In the following reaction

$2A(g) + B(g) \rightleftharpoons 3C(g) + D(g)$ two moles each of A and B are initially taken in a one-litre flask. What will be the $[A] - [D]$?

Q.25 What of the following effect will occur when CN^- is added in the reaction, $HCN(aq) \rightleftharpoons H^+(aq) + CN^-(aq)$, at equilibrium?

Q.26 In the reaction,

$CH_3COOH(l) + C_2H_5OH(l) \rightleftharpoons H_2O(l)$, the concentration of product can be increased when?

Q.27 What happens when pressure is increased in the equilibrium system, $Ice \rightleftharpoons Water$?

Q.28 A vessel at 1000 K contains CO_2 with a pressures of 0.5 atmosphere. Some of the CO_2 in converted into CO on addition on graphite. Calculate the value of K, if total pressure at equilibrium is 0.8 atmosphere.

Q.29 At a certain temperature equilibrium constant (K_c) is 16 for the reaction: $SO_2(g) + NO_2(g) \rightleftharpoons SO_3(g) + NO(g)$. If we take 1 mole of each of the four gases in 1 litre vessel, what is equilibrium concentration of NO and NO_2 ?

Q.30 For the formation of ammonia the equilibrium constant data at 673 K and 773 K respectively are 1.64×10^{-4} and 1.44×10^{-5} respectively. Calculate heat of the reaction. Given $R = 8.314 \text{ JK}^{-1} \text{ mol}^{-1}$.

Q.31 K_c for the reaction $N_2O_4 \rightleftharpoons 2NH_2$ in chloroform at 291 K is 1.14. Calculate the free energy change of the reaction when the concentration of the two gases are 0.5 mol dm^{-3} each at the same temperature. ($R = 0.082 \text{ lit atm K}^{-1} \text{ mol}^{-1}$)

Q.32 Calculate the pressure of CO_2 gas at 700K in the heterogeneous equilibrium reaction

$CaCO_3(s) \rightleftharpoons CaO(s) + CO_2(g)$ if ΔG° for this reaction is $130.2 \text{ kJ mol}^{-1}$.

Q.13 28 g of N_2 and 6 g of H_2 were kept at C in 1 L vessel, the equilibrium mixture contained 27.54 g of NH_3 . The approximate value of K_c for the above reaction can be (in $\text{mol}^{-2} \text{L}^2$):

- (A) 25 (B) 50 (C) 75 (D) 100

Q.14 The equilibrium concentration of X, Y and XY_2 are 4, 2 and 2 moles respectively for the equilibrium, $2X + Y \rightleftharpoons XY_2$. The value of K_c is:

- (A) 0.625 (B) 0.0625 (C) 0.00625 (D) 6.25

Q.15 An amount of solid NH_4HS is placed in a flask already containing ammonia gas at a certain temperature and 0.50 atm pressure. Ammonia hydrogen sulphide decomposes to yield NH_3 and H_2S gases in the flask. When the decomposition reaction reaches equilibrium, the total pressure in the flask rises to 0.84 atm. The equilibrium constant for decomposition at this temperature is:

- (A) 0.11 (B) 0.22 (C) 0.33 (D) 0.44

Q.16 A reaction is, $A + B \longrightarrow C + D$, Initially we start with equal concentration of A, and B. At equilibrium we find the moles of C is two times of A. What is the equilibrium constant of the reaction?

- (A) 2 (B) 4 (C) $\frac{1}{2}$ (D) $\frac{1}{4}$

Q.17 9.2 g of $N_2O_4(g)$ is taken in a closed 1 L vessel and heated till the following equilibrium is reached.

$N_2O_4(g) \rightleftharpoons 2NO_2(g)$. At equilibrium, 50% $N_2O_4(g)$ is dissociated. What is the equilibrium constant (in mol/L) (molecular weight of $N_2O_4 = 92$).

- (A) 0.1 (B) 0.2 (C) 0.3 (D) 0.4

Q.18 In Haber process, 30 L of dihydrogen and 30 L of dinitrogen were taken for reaction which yielded only 50% of the expected product. What will be the composition of gaseous mixture under the aforesaid condition in the end?

- (A) 10 L NH_3 , 25 L NH_3 , 15 L
 (B) 20 L NH_3 , 20 L NH_3 , 20 L
 (C) 20 L NH_3 , 25 L NH_3 , 15 L
 (D) 20 L NH_3 , 10 L NH_3 , 30 L

Q.19 3.2 moles of hydrogen iodine were heated in a sealed bulb at 444°C till the equilibrium state was reached. Its degree of dissociation at this temperature was found to be 22%. The number of moles of hydrogen iodine present at equilibrium are:

- (A) 1.876 (B) 2.496 (C) 3.235 (D) 4.126

Q.20 56 g of nitrogen and 8 g hydrogen gas heated in a closed vessel. At equilibrium, 34 g of ammonia are present. The equilibrium number of moles of nitrogen, hydrogen and ammonia are, respectively:

- (A) 1, 1, 2 (B) 1, 2, 2 (C) 2, 1, 1 (D) 2, 2, 1

Q.21 The reaction, $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$ is carried out in a 1 dm^3 vessel and 2 dm^3 vessel respectively. The ratio of the reaction velocities will be:

- (A) 1 : 4 (B) 2 : 4 (C) 1 : 8 (D) 8 : 1

Q.22 When $NaNO_3$ is heated in a closed vessel, O_2 is liberated and $NaNO_2$ is left behind. At equilibrium:

- (i) Addition of $NaNO_3$ favours forward reaction
 (ii) Addition of $NaNO_2$ favours backward reaction
 (iii) Increasing pressure favours reverse reaction
 (iv) Increasing temperature favours forward reaction

- (A) (i), (ii), (iii) (B) (ii), (iii), (iv)
 (C) (i), (iii), (iv) (D) (i), (ii), (iii), (iv)

Q.23 5 moles of SO_2 and 5 moles of O_2 are allowed to react to form SO_3 in a closed vessel. At the equilibrium stage 60% of SO_2 is used up. The total number of moles of SO_2 , O_2 and SO_3 in the vessel now is:

- (A) 8.5 (B) 9.5 (C) 10 (D) 10.5

Q.24 K for the synthesis of HI is 50. K for the dissociation of HI is:

- (A) 0.2 (B) 0.02 (C) 0.4 (D) 0.04

Q.25 For which of the following reactions $K_p = K_c$?

- (A) $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$
 (B) $H_2(g) + Cl_2(g) \rightleftharpoons 2HCl(g)$
 (C) $2NOCl(g) \rightleftharpoons 2NO(g) + Cl_2(g)$
 (D) $N_2O_4(g) \rightleftharpoons 2NO_2(g)$

Q.26 In which of the following reaction $K_p > K_c$

- (A) $N_2 + 3H_2 \rightleftharpoons 2NH_3$ (B) $H_2 + I_2 \rightleftharpoons 2HI$
 (C) $2SO_3 \rightleftharpoons O_2 + 2SO_2$ (D) $PCl_3 + Cl_2 \rightleftharpoons PCl_5$

Q.27 The equilibrium constant (K_p) for the reaction, $PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$ is 16. If the volume of the container is reduced to one half its original volume, the value of K_p for the reaction at the same temperature will be:

- (A) 8 (B) 16 (C) 32 (D) 64

Q.28 For the reversible reaction, $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$ at $500^\circ C$, the value of K_p is 1.44×10^{-5} when partial pressure is measured in atmospheres. The corresponding value of K_c with concentration in mol L^{-1} , is:

- (A) $\frac{1.44 \times 10^{-5}}{(0.082 \times 773)^{-2}}$ (B) $\frac{1.44 \times 10^{-5}}{(0.082 \times 773)^2}$
 (C) $\frac{1.44 \times 10^{-5}}{(8.314 \times 773)^{-2}}$ (D) $\frac{1.44 \times 10^{-5}}{(0.082 \times 500)^{-2}}$

Q.29 A chemical reaction is catalysed by a catalyst X. Hence, X:

- (A) Reduce enthalpy of the reaction
 (B) Decreases rates constant of the reaction
 (C) Increases rate constant of the reaction
 (D) Does not affect equilibrium constant of reaction

Q.30 Phosphorus pentachloride dissociate as follows, in a closed reaction vessel, $PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$

If total pressure at equilibrium of the reaction mixture is P and degree of dissociation of is x, the partial pressure of will be:

- (A) $\left(\frac{x}{x+1}\right)P$ (B) $\left(\frac{2x}{1-x}\right)P$ (C) $\left(\frac{x}{x-1}\right)P$ (D) $\left(\frac{x}{1-x}\right)P$

Q.31 The equilibrium constant for the reaction

$SO_3(g) \rightleftharpoons SO_2(g) + \frac{1}{2}O_2(g)$. Is $K_c = 4.9 \times 10^{-2}$. The value of K_c for the reaction.

$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$ will be:

- (A) 416 (B) 2.40×10^{-3} (C) 9.8×10^{-2} (D) 4.9×10^{-2}

Previous Years' Questions

Q.1 For the reaction $CO(g) + H_2O(g) \rightleftharpoons CO_2(g) + H_2(g)$

At a given temperature, the equilibrium amount of $CO_2(g)$ can be increased by **[1998]**

- (A) Adding a suitable catalyst
 (B) Adding an inert gas
 (C) Decreasing the volume of the container
 (D) Increasing the amount $CO(g)$

Q.2 At constant temperature, the equilibrium constant (K_p) for the decomposition reaction $N_2O_4 \rightleftharpoons 2NO_2$ is expressed by

$K_p = \frac{(4x^2P)}{(1-x^2)}$, where P = pressure, x = extent of

decomposing. Which one of the following statements is true **[2001]**

- (A) K_p increases with increase of P
 (B) K_p increases with increases of x
 (C) K_p increases with increases of x
 (D) K_p remains constant with change in P & x

Q.3 Ammonia under a pressure of 15 atm at $27^\circ C$ is heated to $437^\circ C$ in a closed vessel in the presence of a catalyst. Under the conditions, NH_3 is partially decomposed according to the equation, $2NH_3 \rightleftharpoons N_2 + 3H_2$. The vessel is such that the volume remains effectively constant whereas pressure increases to 50 atm. Calculate the percentage of NH_3 actually decomposed. **[2001]**

- (A) 65% (B) 61.3% (C) 62.5% (D) 64%

Q.4 The partial pressure of CH_3OH , CO and H_2 in the equilibrium mixture of the reaction $CO + 2H_2 \rightleftharpoons CH_3OH$ at $427^\circ C$ are 2.0, 1.0 and 0.1 atm respectively. The value of K_p for the decomposition of CH_3OH to CO and H_2 is **[1999]**

- (A) $1 \times 10^2 \text{ atm}$ (B) $2 \times 10^2 \text{ atm}^{-1}$
 (C) 50 atm^2 (D) $5 \times 10^{-3} \text{ atm}^2$

Q.5 For the gas phase reaction $C_2H_4 + H_2 \rightleftharpoons C_2H_6$. Carried out in a vessel, the equilibrium concentration of C_2H_4 can be increased by **[1984]**

- (A) Increasing the temperature
 (B) Decreasing the pressure

- (C) Removing some H_2
 (D) Adding some C_2H_6

Q.6 When $NaNO_3$ is heated in a closed vessel, oxygen is liberated and $NaNO_2$ is left behind. At equilibrium

[1986]

- (A) Addition of $NaNO_2$ favours reverse reaction.
 (B) Addition of $NaNO_3$ favours forward reaction.
 (C) Increasing temperature favours forward reaction
 (D) Increasing pressure favours reverse reaction

Q.7 For the reaction $PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$.

The forward reaction at constant temperature is favored by

[1991]

- (A) Introducing an inert gas at constant volume
 (B) Introducing chlorine gas at constant volume
 (C) Introducing an inert gas at constant pressure
 (D) Increasing the volume of the container
 (E) Introducing PCl_5 at constant volume

Q.8 When two reactants, A and B are mixed to give products, C and D, the reaction quotient, (Q) at the initial stages of the reaction

[2000]

- (A) Is zero (B) Decreases with time
 (C) Is independent of time (D) Increases with time

Each of the questions given below consists of two statements, an assertion (Assertion) and reason (Reason). Select the number corresponding to the appropriate alternative as follows


- (A) If both assertion and reason are true and reason is the correct explanation of assertion.
 (B) If both assertion and reason are true and reason is not the correct explanation of assertion.
 (C) If assertion is true but reason is false.
 (D) If assertion is false but reason is true.

Q.9 Assertion: Effect of temperature on K_c or K_p depends on enthalpy change.

[1993]

Reason: Increase in temperature shifts the equilibrium in exothermic direction and decrease in temperature shifts the equilibrium position in endothermic direction.

Q.10 Assertion: For a gaseous reaction,



[1996]

Reason: Concentration of gaseous reactant is taken to be unity.

Q.11 Assertion: Ice \rightleftharpoons water, if pressure is applied water will evaporate.

[1986]

Reason: Increases of pressure pushes the equilibrium towards the side in which number of gaseous mole decreases.

Q.12 For the following three reactions a, b and c, equilibrium constants are given:

[2008]

- a. $CO(g) + H_2O(g) \rightleftharpoons CO_2(g) + H_2(g); K_1$
 b. $CH_4(g) + H_2O(g) \rightleftharpoons CO(g) + 3H_2(g); K_2$
 c. $CH_4(g) + 2H_2O(g) \rightleftharpoons CO_2(g) + 4H_2(g); K_3$

Which of the following relations is correct?

- (A) $K_1 \sqrt{K_2} = K_3$ (B) $K_2 K_3 = K_1$
 (C) $K_3 = K_1 \cdot K_2$ (D) $K_3 \cdot K_2^3 = K_1^2$

Q.13 The equilibrium constant (K_c) for the reaction at temperature T is $N_2(g) + O_2(g) \rightarrow 2NO(g)$. The value of K_c for the reaction, $NO(g) \rightarrow \frac{1}{2} N_2(g) + \frac{1}{2} O_2(g)$ at the same temperature is:

[2012]

- (A) 0.02 (B) 25×10^2 (C) 4×10^{-4} (D) 50.0

Q.14 For the reaction $SO_2(g) + \frac{1}{2} O_2(g) \rightleftharpoons SO_3(g)$, if $K_p = K_c (RT)^x$ where the symbols have usual meaning then the value of x is (assuming ideality)

[2014]

- (A) -1 (B) $-\frac{1}{2}$ (C) $\frac{1}{2}$ (D) 1

Q.15 The standard Gibbs energy change at 300 K for the reaction $2A \rightleftharpoons B + C$ is 2494.2 J. At a given time, the composition of the reaction mixture is $[A] = \frac{1}{2}, [B] = 2$ and $[C] = \frac{1}{2}$. The reaction proceeds in the: [R = 8.314 J/K/mol, e = 2.718]

[2015]

- (A) Forward direction because $Q > K_c$
 (B) Reverse direction because $Q > K_c$
 (C) Forward direction because $Q < K_c$
 (D) Reverse direction because $Q < K_c$

JEE Advanced/Boards

Exercise 1

Q.1 The equilibrium constant K_p for the reaction, $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{SO}_3(\text{g})$ is 900 atm^{-1} at 800 K . A mixture containing SO_3 and O_2 having initial partial pressure of 1 and 2 atm respectively is heated at constant volume to equilibrate. Calculate the partial pressure of each gas at 800 K .

Q.2 What is the concentration of CO in equilibrium at 25°C in a sample of a gas originally containing 1.00 mol L^{-1} of CO_2 ? For the dissociation of CO_2 at 25°C , $K_c = 2.96 \times 10^{-92}$.

Q.3 Ammonia is heated at 15 atm from 27°C to 347°C assuming volume constant. The new pressure becomes 50 atm at equilibrium. Calculate % of NH_3 actually decomposed.

Q.4 Calculate the percent dissociation of $\text{H}_2\text{S}(\text{g})$ if 0.1 mole of H_2S is kept in 0.4 litre vessel at 1000 K for the reaction, $2\text{H}_2\text{S}(\text{g}) \rightleftharpoons 2\text{H}_2(\text{g}) + \text{S}_2(\text{g})$

The value of K_c is 1.0×10^{-6}

Q.5 The vapour density (hydrogen = 1) of a mixture containing NO_2 and N_2O_4 is 38.3 at 26.7°C . Calculate the number of moles of NO_2 in 100 grams of the mixture.


Q.6 At temperature T , the compound $\text{AB}_2(\text{g})$ dissociated according to the reaction, $2\text{AB}_2(\text{g}) \rightleftharpoons 2\text{AB}(\text{g}) + \text{B}_2(\text{g})$.

With a degree of dissociation, x , which is small compared with unity. Deduce the expression for x in terms of the equilibrium constant, K_p and the total pressure, P .


Q.7 At 25°C and one atmospheric pressure, the partial pressures in an equilibrium mixture of N_2O_4 and NO_2 are 0.7 and 0.3 atmosphere, respectively. Calculate the partial pressures of these gases when they are in equilibrium at 25°C and at a total pressure of 10 atmospheres.

Q.8 At 450°C , the equilibrium constant, K_p , for the reaction, $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$. Was found to be 1.6×10^{-5} at a pressure of 200 atm. If N_2 and H_2 are taken in 1 : 3 ratio what is % of NH_3 formed at this temperature?

Q.9 A mixture of SO_2 and O_2 at 1 atmosphere in the ratio of 2 : 1 is passed through a catalyst at 1170°C for attainment of equilibrium. The exit gas is found to contain 87% SO_3 by volume. Calculate K_p for the reaction,




Q.10 At 627°C and one atmosphere SO_3 is partially dissociated into SO_2 and O_2 by the reaction.



The density of the equilibrium mixture is 0.925 g L^{-1} . What is the degree of dissociation?

Q.11 When limestone is heated, quicklime is formed according to the equation.



The experiment was carried out in the temperature range $800 - 900^\circ\text{C}$. Equilibrium constant K_p follows the relation, $\log K_p = 7.282 - 8500 / T$

Where, T is temperature in Kelvin. At what temperature the decomposition will give $\text{CO}_2(\text{g})$ at 1 atm?

Q.12 In the following equilibrium $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$

When 5 moles of each is taken and the temperature is kept at 298 K , the total pressure was found to be 20 bar.

Given: $\Delta G_f^\circ \text{N}_2\text{O}_4 = 100 \text{ kJ}$, $\Delta G_f^\circ \text{NO}_2 = 50 \text{ kJ}$

(i) Find ΔG of the reaction at 298 K .


(ii) Find the direction of the reaction

Q.13 Equilibrium constant for the reaction of iodine with propane according to the following was determined.

Some results obtained at 545 K were as given ahead:

Initial Pressure (mm Hg)			Equilibrium Pressure (mm)	
I_2	C_3H_6	HI	HI	$\text{C}_3\text{H}_5\text{I}$
23.9	505.8	0	1.80	1.80
16.1	355.3	1.62	2.27	0.645

Calculate equilibrium constant according to the following equation,




Q.14 The equilibrium constant K_p for the reaction, $N_2 + 3H_2 \rightleftharpoons 2NH_3$ is 1.64×10^{-4} at 400°C and 0.144×10^{-4} at 500°C . Calculate the mean heat of formation of 1 mole of NH_3 from its elements in this temperature range.

Q.15 When 3.06 g of solid NH_4HS is introduced into a two litre evacuated flask at 27°C , 30% of the solid decomposes into gaseous ammonia and hydrogen sulphide.

- (i) Calculate K_c and K_p for the reaction at 27°C .
 (ii) What would happen to the equilibrium when more solid is introduced into the flask?

Q.16 At 540 K, 0.10 mole of PCl_5 are heated in a 8 litre flask. The pressure of equilibrium mixture is found to be 1.0 atm. Calculate K_p and K_c for the reaction.

Q.17 Density of equilibrium mixture of N_2O_4 and NO_2 at 1 atm and 384 K is 1.84 g dm^{-3} . Calculate the equilibrium constant of the reaction.



Q.18 $2NOBr(g) \longrightarrow 2NO(g) + Br_2(g)$. If nitrosyl bromide ($NOBr$) is 33.33% dissociation at 25°C and a total pressure of 0.28 atm. Calculate K_p for the dissociation at this temperature.

Q.19 At 30°C , the following equilibrium is established: $H_2(g) + S(g) \longrightarrow H_2S(g)$, $K_p = 6.8 \times 10^{-2}$. If 0.2 mol of hydrogen and 1.0 mol of sulphur are heated to 90°C in a 1.0 litre vessel, what will be the partial pressure of H_2S at equilibrium?

Q.20 A mixture of 2 moles of CH_4 and 34 gms of H_2S was placed in an evacuated container, which was then heated to and maintained at 727°C . When equilibrium was established in the gaseous reaction $CH_4 + 2H_2S \rightarrow CS_2 + 4H_2$ the total pressure in the container was 0.92 atm. & the partial pressure of hydrogen was 0.2 atm. What was the volume of the container?

Q.21 At 1200°C , the following equilibrium is established between chlorine atoms & molecules:



The composition of the equilibrium mixture may be determined by measuring the rate of effusion the mixture through a pin hole. It is found that at 1200°C and 1 atm pressure the mixture effuses 1.16 times as fast as krypton effuses under the same condition. Calculate the equilibrium constant K_c .

Q.22 SO_3 decomposes at a temperature of 1000 K and at a total pressure of 1.642 atm. At equilibrium, the density of mixture is 1.28 g/l in a vessel. Find the degree of dissociation of SO_3 for $SO_3(g) \longrightarrow SO_2(g) + 1/2O_2(g)$.

Q.23 Consider the equilibrium: $P(g) + 2Q(g) \rightleftharpoons R(g)$. When the reaction is carried out at a certain temperature, the equilibrium concentration of P and Q are 3M and 4M respectively. When the volume of the vessel is doubled and the equilibrium is allowed to be re-established, the concentration of Q is found to be 3M. Find (i) K_c , (ii) concentration of R at two equilibrium stages.

Exercise 2

Single Correct Choice Type

Q.1 The reaction which proceeds in the forward direction is:

- (A) $Fe_2O_3 + 6HCl = 2FeCl_3 + 3H_2O$
 (B) $SnCl_4 + Hg_2Cl_2 = SnCl_2 + 2HgCl_2$
 (C) $NH_3 + H_2O + NaCl = NH_4Cl + NaOH$
 (D) $2CuI + I_2 + 4K^+ = 2Cu^{2+} + 3KI$

Q.2 In $K_p = K_c [RT]^{\Delta n}$, Δn may have:

- (A) +ve values (B) -ve values
 (C) Integer or fractional values (D) Either of the above

Q.3 For reaction, $PCl_3(g) + Cl_2(g) \rightleftharpoons PCl_5(g)$, the value of K_c at 250°C is 26 mol litre⁻¹. The value of K_p at this temperature will be:

- (A) 0.61 atm⁻¹ (B) 0.57 atm⁻¹
 (C) 0.83 atm⁻¹ (D) 0.46 atm⁻¹


Q.4 For the reversible reaction, $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$ at 500°C , the value of K_p is 1.44×10^{-5} when partial pressure is measured in the atmosphere. The corresponding value of K_c with concentration in mol litre⁻¹, is:

- (A) $1.44 \times 10^{-5} / (0.082 \times 500)^{-2}$
 (B) $1.44 \times 10^{-5} / (8.314 \times 773)^{-2}$
 (C) $1.44 \times 10^{-5} / (0.082 \times 773)^{-2}$
 (D) $1.44 \times 10^{-5} / (0.082 \times 773)^{-2}$

Q.5 A cylinder fitted with a movable piston contains liquid water in equilibrium with water vapour at 25°C. Which operation result in a decrease in the equilibrium vapour pressure?

- (A) Moving the piston downward a short distance
- (B) Removing a small amount of vapour
- (C) Removing a small amount of the liquid water
- (D) Dissolving salt in the water

Q.6 The volume of the reaction vessel containing an equilibrium mixture in the reaction,



Is increases. When equilibrium is reestablished:

- (A) The amount of $\text{SO}_2(\text{g})$ will decrease
- (B) The amount of $\text{SO}_2\text{Cl}_2(\text{g})$ will decrease
- (C) The amount of $\text{Cl}_2(\text{g})$ will increase
- (D) The amount of $\text{Cl}_2(\text{g})$ will remain unchanged.

Q.7 For the equilibrium $\text{CaCO}_3(\text{s}) \rightleftharpoons \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$, which of the following expression is correct?

- (A) $K_p = [\text{CaO}][\text{CO}_2] / [\text{CaCO}_3]$
- (B) $K_p = (P_{\text{CaO}} \times P_{\text{CO}_2}) / P_{\text{CaCO}_3}$
- (C) $K_p = P_{\text{CO}_2}$
- (D) $K_p = (P_{\text{CaO}} + P_{\text{CO}_2}) / P_{\text{CaCO}_3}$

Q.8 Just before a reversible reaction attains equilibrium it is found that:

- (A) The velocity of both forward reaction and backward reaction is also increasing
- (B) The velocity of the forward reaction is decreasing and that of backward reaction is increasing
- (C) The velocity of both forward and backward reaction is decreasing
- (D) All of the above

Q.9 Densities of diamond and graphite are 3.5 and 2.3 g/mL respectively. Increase of pressure on the equilibrium $\text{C}_{(\text{diamond})} \rightleftharpoons \text{C}_{(\text{graphite})}$:

- (A) Favours backward reaction
- (B) Favours forward reaction
- (C) Have no effect
- (D) Increases the reaction rate

Q.10 For the reaction, $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$ in a vessel, after the addition of equal numbers of mole of N_2 and H_2 , equilibrium state is formed. Which of the following is correct?

- (A) $[\text{H}_2] = [\text{N}_2]$
- (B) $[\text{H}_2] < [\text{N}_2]$
- (C) $[\text{H}_2] > [\text{N}_2]$
- (D) $[\text{H}_2] > [\text{NH}_3]$

Q.11 A characteristic feature of reversible reaction is that:

- (A) They never proceed to completion
- (B) They proceed to completion
- (C) They are not complete unless the reactants are removed from sphere of reaction mixture
- (D) None of the above

Q.12 An example of reversible reaction is:

- (A) $\text{AgNO}_3(\text{aq}) + \text{HCl}(\text{aq}) \longrightarrow \text{AgCl}(\text{s}) + \text{HNO}_3(\text{aq})$
- (B) $2\text{Na} + 2\text{H}_2\text{O} \longrightarrow 2\text{NaOH} + \text{H}_2$
- (C) $\text{NaOH} + \text{CH}_3\text{COOH} \longrightarrow \text{CH}_3\text{COONa} + \text{H}_2\text{O}$
- (D) $\text{Pb}(\text{NO}_3)_2 + 2\text{NaI} \longrightarrow \text{PbI}_2 + 2\text{NaNO}_3$

Q.13 The correct relation between K_p and K_c for the reaction $a\text{X} + b\text{Y} \rightleftharpoons b\text{Z} + a\text{W}$ is:

- (A) $K_p = K_c [\text{Rt}]^{a+b}$
- (B) $K_p = \frac{K_c}{(a+b)^2}$
- (C) $K_p = K_c [\text{RT}]$
- (D) $K_p = K_c$

Q.14 The reaction that proceeds in the forward direction is:

- (A) $\text{SnCl}_4 + \text{Hg}_2\text{Cl}_2 \longrightarrow \text{SnCl}_2 + 2\text{HgCl}_2$
- (B) $\text{NH}_4\text{Cl} + \text{NaOH} \longrightarrow \text{H}_2\text{O} + \text{NH}_3 + \text{NaCl}$
- (C) $\text{Mn}^{2+} + 2\text{H}_2\text{O} + \text{Cl}_2 \longrightarrow \text{MnO}_2 + 4\text{H}^+ + 2\text{Cl}^-$
- (D) $\text{S}_4\text{O}_6^{2-} + 2\text{I}^- \longrightarrow 2\text{S}_2\text{O}_3^{2-} + \text{I}_2$

Q.15 Which information can be obtained from Le Chatelier's principle?

- (A) Shift in equilibrium position on changing P, T and concentration
- (B) Dissociation constant of a weak acid
- (C) Energy change in a reaction
- (D) Equilibrium constant of a chemical reaction

Q.16 The values of K_{p_1} and K_{p_2} for the reaction:



Are in the ratio 9 : 1. Assuming degree of dissociation of X and A and be same, the dissociation pressure at equilibrium (i) and (ii) are in the ratio:

- (A) 36 : 1 (B) 1 : 1 (C) 3 : 1 (D) 1 : 9

Q.17 For the equilibrium, $PCl_5 \rightleftharpoons PCl_3 + Cl_2$

$K_c = \alpha^2 / (1 - \alpha)V$, temperature remaining constant:

- (A) K_c will increase with the increase in volume
 (B) K_c will increase with decrease in volume
 (C) K_c will not change with the change in volume
 (D) K_c may increase or decrease with the change in volume depending upon its numerical value.

Q.18 Consider the following equilibrium in a closed container: $N_2O_4(g) \rightleftharpoons 2NO_2(g)$

At a fixed temperature, the volume of the reaction container is halved. For this change, which of the given statements holds true regarding the equilibrium constant (K_p) and degree of dissociation (α)?

- (A) Neither K_p nor α changes
 (B) Both K_p and α change
 (C) K_p changes, but α does not change
 (D) K_p does not change, but α changes


Q.19 At constant temperature, the equilibrium constant (K_p) for the decomposition reaction $N_2O_4 \rightleftharpoons 2NO_2$ is

expressed by, $K_p = \frac{(4x^2P)}{(1-x^2)}$

Where, P = pressure, x = extent of decomposition. Which of the following statements is true?

- (A) K_p increases with increase of P
 (B) K_p increases with increases of x
 (C) K_p increases with decrease of x
 (D) K_p remains constant with change in P and x decreases with pressure

Q.20 For the reaction in equilibrium,



The concentrations, of N_2O_4 and NO_2 at equilibrium are 4.8×10^{-2} and 1.2×10^{-2} mol L⁻¹ respectively. The value of K_c for this reaction is:

- (A) 3×10^{-3} mol L⁻¹ (B) 3×10^3 mol L⁻¹
 (C) 3.3×10^2 mol L⁻¹ (D) 3×10^{-1} mol L⁻¹

Q.21 1.1 mole of A are mixed with 2.2 mole of B and the mixture is then kept in one litre flask till the equilibrium is attained $A + 2B \rightleftharpoons 2C + D$. At the equilibrium, 0.2 mole of C are formed. The equilibrium constant of the reaction is:

- (A) 0.001 (B) 0.002 (C) 0.003 (D) 0.004


Q.22 For the reactions, $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$ at 720 K, the value of equilibrium constant is 50, when equilibrium concentration of both H_2 and I_2 is 0.5 M. K_p under the same conditions will be:

- (A) 0.02 (B) 0.2 (C) 50 (D) 50 RT

Q.23 The equilibrium constant for a reaction is 1×10^{20} at 300 K. The standard Gibbs energy change for this reaction is:

- (A) - 115 kJ (B) + 115 kJ (C) + 166 kJ (D) - 166 kJ

Q.24 The equilibrium constant for the reaction:



- (A) $K_c = \frac{1}{[O_2]^5}$ (B) $K_c = [O_2]^5$
 (C) $K_c = \frac{[P_4O_{10}]}{5[P_4][O_2]}$ (D) $K_c = \frac{[P_4O_{10}]}{5[P_4][O_2]^5}$

Q.25 Eight mole of a gas AB_3 attain equilibrium in a closed container of volume 1 dm³ as, $2AB_3 \rightleftharpoons A_2(g) + 3B_2(g)$. If at equilibrium 2 mole of A_2 are present then, equilibrium constant is:

- (A) 72 mol² L⁻² (B) 36 mol² L⁻²
 (C) 3 mol² L⁻² (D) 27 mol² L⁻²

Q.26 The equilibrium constant for the reaction, $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}(\text{g})$ is 4×10^{-4} at 2000 K. In presence of a catalyst the equilibrium is attained 10 times faster. Therefore, the equilibrium constant, in presence of the catalyst, at 2000 K is:

- (A) 40×10^{-4}
 (B) 4×10^{-4}
 (C) 4×10^{-3}
 (D) Difficult to compute without more data

Q.27 The reaction, $\text{A} + 2\text{B} \rightleftharpoons 2\text{C} + \text{D}$ was studied using an initial concentration of B which was 1.5 times that of A. But the equilibrium concentrations of A and C were found to be equal. Then the K_c for the equilibrium is:

- (A) 4 (B) 8 (C) 6 (D) 0.32

Multiple Correct Choice Type

Q.28 The yield of product in the reaction $2\text{A}(\text{g}) + \text{B}(\text{g}) \rightleftharpoons 2\text{C}(\text{g}) + \text{QKJ}$

Would be lower at:

- (A) Low temperature and low pressure
 (B) High temperature and high pressure
 (C) Low temperature and to high pressure
 (D) High temperature and low pressure

Q.29 What is the effect of the reduction of the volume of the system for the equilibrium $2\text{C}(\text{s}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{CO}(\text{g})$?


- (A) The equilibrium will be shifted to the left by the increased pressure caused by the reduction in volume
 (B) The equilibrium will be shifted to the right by the decreased pressure by the reduction in volume.
 (C) The equilibrium will be shifted to the left by the increased pressure caused by the increased in volume.
 (D) The equilibrium will be shifted to the right by the increased pressured caused by the reduction in volume.

Q.30 For the reaction $\text{PCl}_5(\text{g}) \rightleftharpoons \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$, the forward reaction at constant temperature is favoured by

- (A) Introducing an inert gas at constant volume
 (B) Introducing chlorine gas at constant volume
 (C) Introducing an inert gas at constant pressure
 (D) Introducing PCl_5 at constant volume.

Comprehension Type

Paragraph 1: 10 moles of NH_3 is heated at 15 atm from 27°C to 347°C assuming volume constant. The pressure at equilibrium is found to be 50 atm. The equilibrium constant for dissociation of



Can be written as; $K_p = \frac{P_{\text{N}_2} \times (P_{\text{H}_2})^3}{(P_{\text{NH}_3})^2} (\text{atm})^2$

Q.31 The degree of dissociation of is:

- (A) 61.3% (B) 20% (C) 48% (D) None of these

Q.32 The equilibrium constant K_c for the reaction is:

- (A) 7.08×10^2 (B) 3.06×10^2
 (C) 7.6×10^2 (D) 1.53×10^3

Q.33 The volume of container in which gas is heated is:

- (A) 16.42 litre (B) 8.21 litre (C) 20 litre (D) 15 litre

Paragraph 2: For a reversible reaction at a certain temperature when it is at equilibrium has been attained whether physical or chemical, a change in certain variables might change the state of equilibrium. These variables includes pressure, volume, concentration and temperature. Due to these changes, a system under equilibrium changes its state in such a manner, i.e., the equilibrium moves in forward direction or backward direction, so that the effect of change is annulled. For a gaseous phase endothermic decomposition of phosphorus pentachloride, can be made spontaneous by increasing concentration of PCl_5 , lowering the pressure and increasing temperature of the system.

Q.34 Which of the following reactions proceed in forward direction with increase in temperature are:

- (A) $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightleftharpoons 2\text{HI}(\text{g}) + 3000 \text{ cal}$
 (B) $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}(\text{g}) - 43200 \text{ cal}$
 (C) $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{HN}_3(\text{g}) + 22400 \text{ cal}$
 (D) $\text{C}(\text{s}) + \text{O}_2(\text{g}) \longrightarrow \text{CO}_2(\text{g}) + 94300 \text{ cal}$

Q.35 The equilibrium, solid \rightleftharpoons liquid \rightleftharpoons gas, will shift in forward direction when:

- (A) Temperature is raised (B) Temperature is constant
(C) Temperature is lowered (D) Pressure is increases

Q.36 The change in standard Gibbs energy for a reaction at equilibrium, e.g., $\text{PCl}_5(\text{g}) \rightleftharpoons \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$, on addition of an inert gas at constant pressure and then at constant volume respectively are:

- (A) Decreases, no change (B) Increases, no change
(C) No change, no change (D) No change, decreases

Previous Years' Questions

Q.1 For the reaction $\text{CO}(\text{g}) + 2\text{H}_2(\text{g}) \rightleftharpoons \text{CH}_3\text{OH}(\text{g})$, true condition is **[1992]**

- (A) $K_p = K_c$ (B) $K_p > K_c$
(C) $K_p < K_c$ (D) $K_c = 0$ but $K_p \neq 0$

Q.2 Which of the following is not favorable for SO_3 formation $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{SO}_3(\text{g}); \Delta H = -45.0 \text{ kcal}$ **[1997]**

- (A) High pressure
(B) High temperature
(C) Decreasing SO_3 concentration
(D) Increasing reactant concentration

Q.3 The formation of NO_2 in the reaction $2\text{NO} + \text{O}_2 \rightleftharpoons 2\text{NO}_2 + \text{heat}$ is favoured by **[1998]**

- (A) Low pressure
(B) High pressure
(C) Low temperature
(D) Reduction in the mass of reactant

Q.4 For the gas phase reaction $\text{C}_2\text{H}_4 + \text{H}_2 \rightleftharpoons \text{C}_2\text{H}_6$ ($\Delta H = -32.7 \text{ kcal}$). Carried out in a vessel, the equilibrium concentration of C_2H_4 can be increased by **[1984]**

- (A) Increasing the temperature
(B) Decreasing the pressure
(C) Removing some H_2
(D) Adding some C_2H_6

Q.5 The equilibrium $\text{SO}_2\text{Cl}_2(\text{g}) \rightleftharpoons \text{SO}_2(\text{g}) + \text{Cl}_2(\text{g})$ is attained at 25°C in a closed container and an inert gas helium is introduced. Which of the following statements are correct? **[1989]**

- (A) Concentration of SO_2 , Cl_2 and SO_2Cl_2 change
(B) More chlorine is formed
(C) Concentration of SO_2 is reduced
(D) All the above are incorrect

Q.6 For the chemical reaction $3\text{X}(\text{g}) + \text{Y}(\text{g}) \rightleftharpoons \text{X}_3\text{Y}(\text{g})$, the amount of X_3Y at equilibrium is affected by **[2009]**

- (A) Temperature and pressure
(B) Temperature only
(C) Pressure only
(D) Temperature, pressure and catalyst

Q.7 The equilibrium $\text{SO}_2\text{Cl}_2(\text{g}) \rightleftharpoons \text{SO}_2(\text{g}) + \text{Cl}_2(\text{g})$ is attained at 25°C in a closed container and an inert gas helium is introduced which of the following statement is correct. **[1989]**

- (A) More chlorine is formed
(B) Concentration of SO_2 is reduced
(C) More SO_2Cl_2 is formed
(D) Concentration of SO_2Cl_2 , SO_2 and Cl_2 does not

Q.8 In the reaction, $\text{A}_2(\text{g}) + 4\text{B}_2(\text{g}) \rightleftharpoons 2\text{AB}_4(\text{g})$ $\Delta H < 0$ the formation of AB_4 is will be favoured at **[1990]**

- (A) Low temperature, high pressure
(B) Low temperature, low pressure
(C) Low temperature low pressure
(D) High temperature low pressure

Q.9 For the given equilibrium: **[2003]**
 $2\text{SO}_2 + \text{O}_2 \rightleftharpoons 2\text{SO}_3 + \text{heat}$

The equilibrium reaction proceeds in forward direction when

- (A) Oxygen is removed (B) SO_3 is added
(C) Heat is added (D) Oxygen is added

Q.10 Statement-I: For every chemical reaction at equilibrium, standard Gibbs energy of reaction is zero.

Statement-II: At constant temperature and pressure, chemical reactions are spontaneous in the direction of decreasing Gibbs energy. **[2008]**


(A) Statement-I is True, statement-II is True; statement-II is correct explanation for statement-I

(B) Statement-I is True, statement-II is True; statement-II is NOT a correct explanation for statement-I

(C) Statement-I is True, statement-II is False

(D) Statement-I is False, statement-II is True

Q.11 The thermal dissociation equilibrium of (s) is studied under different conditions.



For this equilibrium, the correct statement(s) is(are) [2013]

(A) ΔH is dependent on T

(B) K is independent of the initial amount of CaCO_3

(C) K is dependent on the pressure of CO_2 at a given T

(D) ΔH is independent of the catalyst, if any

Paragraph: Thermal decomposition of gaseous X_2 to gaseous X at 298 K takes place according to the following equation:

The standard reaction Gibbs energy, of this reaction is positive. At the start of the reaction, there is one mole of X_2 and no X. As the reaction proceeds, the number

of moles of X formed is given by β . Thus, $\beta_{\text{equilibrium}}$ is the number of moles of X formed at equilibrium. The reaction is carried out at a constant total pressure of 2 bar. Consider the gases to behave ideally. (Given: $R = 0.083 \text{ L bar K mol}^{-1}$)

Q.12 The equilibrium constant K_p for this reaction at 298 K, in terms of $\beta_{\text{equilibrium}}$, is [2016]

(A) $\frac{8\beta_{\text{equilibrium}}^2}{2 - \beta_{\text{equilibrium}}}$ (B) $\frac{8\beta_{\text{equilibrium}}^2}{4 - \beta_{\text{equilibrium}}^2}$

(C) $\frac{4\beta_{\text{equilibrium}}^2}{2 - \beta_{\text{equilibrium}}}$ (D) $\frac{4\beta_{\text{equilibrium}}^2}{4 - \beta_{\text{equilibrium}}^2}$

Q.13 The INCORRECT statement among the following, for this reaction, is [2016]

(A) Decrease in the total pressure will result in formation of more moles of gaseous X.

(B) At the start of the reaction, dissociation of gaseous X_2 , takes place spontaneously.

(C) $\beta_{\text{equilibrium}} = 0.7$.

(D) $K_c < 1$

MASTERJEE Essential Questions

JEE Main/Boards

Exercise 1

Q.1	Q.7	Q.8
Q.15	Q.17	Q.22
Q.25	Q.29	

Exercise 2

Q.2	Q.6	Q.10
Q.15	Q.17	Q.20
Q.24	Q.29	

Previous Years' Questions

Q.1	Q.3	Q.8
-----	-----	-----

JEE Advanced/Boards

Exercise 1

Q.2	Q.6	Q.8
Q.12	Q.15	Q.23

Exercise 2

Q.10	Q.12	Q.18-20
------	------	---------

Previous Years' Questions

Q.5	Q.6
-----	-----

Answer Key

JEE Main/Boards

Exercise 1

Q.1 At equilibrium: Mole of $I_2 = 0.0075$ moles; Mole of HI = 0.347 moles

Q.2 1:10

Q.3 $K_c = 1.8$

Q.4 $K_p = 1.64 \times 10^{-4}$ atm

Q.5 $K = 1.7 \text{ mol}^2 \text{ l}^{-2}$

Q.6 $\frac{K_p}{K_c} = 1$

Q.7 Partial pressure of HCl = 600 mm

Q.8 $K' = 0.25$

Q.9 $K_c = 0.76$

Q.10 $K_p = 1$ atmosphere

Q.11 $x = 2/5$

Q.12 $P_{\text{NH}_3} = \frac{1}{3}P$

Q.13 $[N_2] = 1.25$

Q.14 $K_p = 3.5 \times 10^{23}$ atm $K_c = 1.4 \times 10^{22} \text{ mol l}^{-1}$

Q.15 0.33

Q.16 $K_2 = \frac{1}{K_1}$

Q.17 1000 times of K_c

Q.18 Remains unchanged

Q.19 Double

Q.20 $K_p = K_c = K_x = 0.1202$

Q.21 3

Q.22 0.8

Q.23 $\frac{x}{a+x}$

Q.24 $[A] - [D] = 2 - 3x$

Q.25 Increase is the negative log value of concentration of H^+

Q.26 H_2SO_4 is used in the reaction, due to absorption of water where the reaction no more remains reversible. Thus, the reaction proceeds in forward direction only and the product is obtained in larger amounts

Q.27 Melting point of ice decreases

Q.28 1.2 atmosphere

Q.29 $NO_2 = 0.4$ mole, $NO = 1.6$ mole

Q.30 -105.216 kJ

Q.31 -19.67 lit atm

Q.32 $P_{CO_2} = K_p = 1.94 \times 10^{-10}$ atm

Q.33 $T = 399$ K.

Exercise 2

Single Correct Choice Type

Q.1 A

Q.2 B

Q.3 B

Q.4 D

Q.5 C

Q.6 A

Q.7 A

Q.8 D

Q.9 B

Q.10 C

Q.11 B

Q.12 A

Q.13 C

Q.14 B

Q.15 A

Q.16 B

Q.17 B

Q.18 A

Q.19 B

Q.20 A

Q.21 D

Q.22 D

Q.23 A

Q.24 B

Q.25 B

Q.26 C

Q.27 B

Q.28 A

Q.29 D

Q.30 A

Q.31 A

Previous Years' Questions

Q.1 D	Q.2 D	Q.3 B	Q.4 D	Q.5 A, B, C, D	Q.6 C, D	Q.7 C, D, E
Q.8 A, D	Q.9 C	Q.10 D	Q.11 E	Q.12 C	Q.13 D	Q.14 B
Q.15 B						

JEE Advanced/Boards

Exercise 1

Q.1 0.977 atm	Q.2 = $3.90 \times 10^{-31} \text{ mol L}^{-1}$	Q.3 61.3	Q.4 2.0	Q.5 0.43 mol NO ₂
Q.6 $x = \left(\frac{2K_p}{P}\right)^{1/3}$	Q.7 $p_{\text{N}_2\text{O}_4} = 8.93 \text{ atm}; p_{\text{NO}_2} = 1.07 \text{ atm}$	Q.8 17.64	Q.9 = $48.24 \text{ atm}^{-1/2}$	
Q.10 34%	Q.11 894.26°C			
Q.12 -5.705 kJ and Negative value shows that reaction will be in forward direction.				
Q.13 2.9×10^{-4}	Q.14 = -12.57 kcal mol ⁻¹	Q.15 0.049	Q.16 1.77 atm	
Q.17 2.09 atm	Q.18 $K_p = 0.01 \text{ atm}$	Q.19 0.379 atm	Q.20 300 L	
Q.21 6.71×10^{-4}	Q.22 $\alpha = 0.5$	Q.23 $K_c = 1/12, [R] = 4 \text{ (initial),} = 1.5 \text{ (final)}$		

Exercise 2

Single Correct Choice Type

Q.1 A	Q.2 D	Q.3 A	Q.4 D	Q.5 D	Q.6 C	Q.7 C
Q.8 B	Q.9 C	Q.10 B	Q.11 A	Q.12 C	Q.13 D	Q.14 B
Q.15 A	Q.16 A	Q.17 C	Q.18 D	Q.19 D	Q.20 A	Q.21 A
Q.22 C	Q.23 A	Q.24 A	Q.25 D	Q.26 B	Q.27 D	

Multiple Correct Choice Type

Q.28 A, B, C	Q.29 A, B, C, D	Q.30 C, D
--------------	-----------------	-----------

Comprehension Type

Paragraph 1: Q.31 A	Q.32 D	Q.33 A
Paragraph 2: Q.34 B	Q.35 A	Q.36 C


Previous Years' Questions

Q.1 C	Q.2 B	Q.3 B, C	Q.4 A, B, C, D	Q.5 D	Q.6 A	Q.7 D
Q.8 A	Q.9 D	Q.10 D	Q.11 A, B, D	Q.12 B	Q.13 C	


Solutions

JEE Main/Boards


Exercise 1



Sol 2:



Sol 3:




Sol 4:

$$K_p = K_c (RT)^{\Delta n} = 0.1768(0.0821 \times 400)^{2-} = 1.64 \times 10^{-4}$$


Sol 5: Let α be the degree of dissociation:

Hence we have:



$$[\text{N}_2] = \frac{1}{2}, [\text{H}_2] = \frac{3}{2}, [\text{NH}_3] = 1$$

$$\therefore K = \frac{[\text{N}_2][\text{H}_2]^3}{[\text{NH}_3]^2} = \frac{\frac{1}{2} \times \frac{3}{2} \times \frac{3}{2} \times \frac{3}{2}}{1 \times 1} = \frac{27}{16} = 1.7 \text{ mol}^2 \text{ l}^{-2}$$





$$K_p = K_c (RT)^{\Delta n}$$

$$\Delta n = 1$$

$$K_c = K_p$$



$$\frac{K_c}{K_p} = 1$$



3/2 moles of Cl_2 will react with 3/2 moles of H_2 to give $\frac{3}{2} \times 2 = 3.0$ moles of HCl . So the total number of moles in the reaction.


$$= \left(2 - \frac{3}{2} \right) + 0 + 3 = 3.5$$

$$\text{Partial pressure of HCl} = \frac{3 \times 700}{3.5} = 600 \text{ mm}$$



$$\therefore \text{NH}_3 \rightleftharpoons \frac{1}{2}\text{N}_2 + \frac{3}{2}\text{H}_2 \rightarrow k^{11} = \sqrt{\frac{1}{16}}$$

Sol 9:




$$K_c = \frac{0.30/3}{\left(\frac{0.70}{3}\right)\left(\frac{1.70}{3}\right)}$$

Sol 10: Total Moles = 2 + 2 + 2 = 6

$$P_{\text{PCl}_3} = \frac{2}{6} \times 3, P_{\text{PCl}_5} = \frac{2}{6} \times 3, P_{\text{Cl}_2} = \frac{2}{6} \times 3$$

$$K_p = \frac{P_{\text{PCl}_3} \times P_{\text{Cl}_2}}{P_{\text{PCl}_5}} = \frac{1 \times 1}{1} = 1 \text{ atmosphere}$$

Sol 11: $\text{H}_2 + \text{I}_2 \rightleftharpoons 2\text{HI}$ $K_c = 9$




$$\begin{array}{cccc} \text{Initial} & 2 & 0 & 0 \\ \text{eq}^m & 2-2x & x & x \end{array}$$

$$\therefore K_c' = \frac{x^2}{(2-2x)^2}; (2-2x)^2 = 9x^2$$


$$2-2x = 3x \therefore x = \frac{2}{5}$$

Sol 12:




$$\therefore P_{\text{NH}_3} = \frac{1}{3} \times P \quad (3 = \text{Total no. of moles})$$


Sol 13:



Sol 14: T = 300 K




$$k_p = 1.7 \times 10^{-12}$$



$$k_p' = \left(\frac{1}{k_p}\right)^2 = \left(\frac{1}{1.7 \times 10^{-12}}\right)^2$$

$$k_p = k_c (RT)^{\Delta n}$$

Sol 15:



$$\begin{array}{ccc} \text{Initial} & \frac{2}{3} & 0 & 0 \end{array}$$

$$\begin{array}{ccc} \text{eq}^m & \frac{2}{3} - \frac{1}{3} & \frac{1}{3} & \frac{1}{3} \end{array}$$

$$\therefore k_c = 0.33$$

Sol 16: $2\text{C} \rightleftharpoons \text{A} + 3\text{B}$

$$k_2 = \frac{1}{k_1} \quad (\because k_1 \Rightarrow \text{A} + 3\text{B} \rightleftharpoons 2\text{C})$$

Sol 17: $\text{P} + \text{Q} \rightleftharpoons \text{R} + \text{S}$

$$K_c = 10^{-2} \quad ; \quad K_f = 10^{-1}$$

$$\therefore K_c = \frac{K_f}{K_b} \Rightarrow K_b = 10^1$$

Sol 18: $\text{A} + 2\text{B} \rightleftharpoons \text{C} + 3\text{D}$


Le-chatlier's principle

Sol 19: Apply law of mass action

Sol 20: In this case


$$\Delta n = 0$$

$$\therefore K_p = K_c = K_x = K$$




Reaction is reversed and multiplied by two.


$$\therefore K = \frac{(8.32)^2}{8.32} = 8.32$$



Reaction is multiplied by 2

$$\therefore K = (8.32)^2 = 69.66$$




Sol 21:**Sol 22:** Vapour density = 57.9

Molar mass = 208.5

$$\alpha = \frac{D-d}{(n-1)d} \Rightarrow \begin{array}{l} D = \text{initial V.D} \\ d = \text{equilibrium V.D.} = 57.9 \end{array}$$

$$D = \frac{M_{\text{PCl}_5}}{2} = \frac{208.5}{2} = 104.25 \quad (n = 2)$$


$$\therefore \alpha = \frac{104.25 - 57.9}{1 \times 57.9} = 0.8$$

Sol 23:

$$\begin{aligned} \therefore \text{Total no. of moles} &= (a-x) + x + x \\ &= a+x \end{aligned}$$


$$\therefore P_{\text{PCl}_5} = \frac{a-x}{a+x}P; \quad P_{\text{PCl}_3} = \frac{x}{a+x}P; \quad P_{\text{Cl}_2} = \frac{x}{a+x}P$$

$$\therefore P_{\text{PCl}_3} \times \frac{1}{P} = \frac{x}{a+x}P \times \frac{1}{P} = \frac{x}{a+x}$$

Sol 24:


$$\therefore [\text{A}] - [\text{D}] = 2 - 2x - x = 2 - 3x$$

Sol 25: What of the following effect will occur when CN^- is added in the reaction, $\text{HCN}(\text{aq}) \rightleftharpoons \text{H}^+(\text{aq}) + \text{CN}^-(\text{aq})$, at equilibrium Increase is the negative log value of concentration of H^+ .

Sol 26:

Le-chatlier's principle

Sol 27: Le-chatlier's principle when pressure is increased in the equilibrium system of water and ice, Melting point of ice decreases.

Sol 28:

Initial Pressure 0.5


Final Pressure $0.5 - x$ $2x$ Total pressure = $0.5 + x$

But the total pressure at equilibrium is 0.8 atm


$$0.5 + x = 0.8$$

$$x = 3 \text{ atm}$$

$$K = \frac{[\text{CO}]^2}{[\text{CO}_2]} = \frac{6^2}{0.3} = 1.2$$

Sol 29:

$$k_c = 16$$



$$k_c = \frac{x^2}{(1-x)^2} \Rightarrow 4 = \frac{x}{1-x}$$

$$4 - 4x = x; \quad 4 = 5x \quad 0$$

$$x = \frac{4}{5} = 0.8$$

$\therefore \text{eq}^m \text{ conc. of NO} = 0.8$

$$\text{NO}_2 = 0.2$$

Sol 30:

$$T_1 = 673\text{k}$$

$$T_2 = 773\text{k}$$

$$k_1 = 1.64 \times 10^{-4}$$

$$k_2 = 1.44 \times 10^{-5}$$

$$R = 8.314 \text{ Jk}^{-1} \text{ mol}^{-1}$$

$$\log \frac{k_2}{k_1} = \frac{\Delta H}{2.303R} \cdot \left(\frac{1}{T_1} - \frac{1}{T_2} \right) \quad (\text{Apply})$$


Sol 31: $\text{N}_2\text{O}_4 \rightleftharpoons 2\text{NO}_2$

$$k_c = 1.14$$

$$\Delta G = ? \quad ; \quad R = 0.082 \text{ lit atm k}^{-1} \text{ mol}^{-1}$$

$$T = 291\text{K}$$


$$\Delta G = -RT \ln k \quad (\text{Apply})$$



$$\Delta G^\circ = 130.2 \text{ k J mol}^{-1}$$

$$T = 700 \text{ k}$$

$$\Delta G^\circ = -RT \ln k \quad (\text{Apply})$$



$$\Delta G^\circ = -20,700 - 11.97T$$

$$\Delta G^\circ = -RT \ln k$$

$$-20,700 - 11.97T = -0.082T \ln k$$

Exercise 2

Single Correct Choice Type

Sol 1: (A) As we increases the concentration of substance, then speed of the reaction increases.

Sol 2: (B) Chemical reaction quantitatively depend on the reactant and product molecule.

Sol 3: (B) In the thermal decomposition of potassium chlorate given as:


$2\text{KClO}_3 \longrightarrow 2\text{KCl} + 3\text{O}_2$, law of mass action cannot be applied.

Sol 4: (D) Those reaction which have more value of K proceeds towards completion.

Sol 5: (C) K_c is a characteristic constant for the given reaction.

Sol 6: (A) K_p is a constant and does not change with pressure.

Sol 7: (A)



Volume of container = 2 L.


$$K_c = \frac{\frac{2 \times 40}{100 \times 2} \times \frac{2 \times 40}{100 \times 2}}{\frac{2 \times 60}{100 \times 2}} = 0.266$$

Sol 8: (D)

$$K_c = \frac{[\text{NO}_2]^2}{[\text{N}_2\text{O}_4]} = \frac{\left[\frac{2 \times 10^{-3}}{2} \right]^2}{\left[\frac{0.2}{2} \right]}$$


$$= \frac{10^{-6}}{10^{-1}} = 10^{-5}$$

Sol 9: (B)



$$K_c = \frac{[\text{C}][\text{D}]}{[\text{A}][\text{B}]} = \frac{2x \cdot 2x}{x \cdot x} = 4$$

Sol 10: (C)



$$\text{Total moles at equilibrium} = (1 - \alpha) + 2\alpha$$

$$= 1 + \alpha$$

Sol 11: (B)

$$K_c = \frac{[\text{HI}]^2}{[\text{H}_2][\text{I}_2]}$$

$$\therefore 64 = \frac{x^2}{0.03 \times 0.03}$$

$$\therefore x^2 = 64 \times 9 \times 10^{-4}$$


$$\text{Or, } x = 8 \times 3 \times 10^{-2}$$

x is the amount of HI at equilibrium amount of I_2 at equilibrium will be:

$$0.30 - 0.24 = 0.06$$

Sol 12: (A) $K_c = \frac{k_f}{k_b}$

$$\therefore k_f = k_c \times k_b = 1.5 \times 7.5 \times 10^{-4} = 1.125 \times 10^{-3}$$

Sol 13: (C)


$$\text{Number of moles of N}_2 = \frac{28}{28} = 1 \text{ mol}$$

$$\text{Number of moles of H}_2 = \frac{6}{2} = 3 \text{ mol}$$

$$\text{Number of moles of NH}_3 = \frac{27.54}{17} = 1.62 \text{ mol}$$

$$\therefore K_c = \frac{[\text{NH}_3]^2}{[\text{N}_2][\text{H}_2]^3} = \frac{[1.62]^2}{[0.19][0.57]^3} = 75$$


$$\text{Sol 14: (B)} \quad K_c = \frac{[\text{YX}_2]}{[\text{X}]^2[\text{Y}]} = \frac{2}{4 \times 4 \times 2} = \frac{1}{16} = 0.0625$$

Sol 15: (A)

$$\text{Total pressure} = 0.5 + 2x = 0.84$$

$$\text{i.e., } x = 0.17$$

$$K_p = P_{\text{NH}_3} \cdot P_{\text{H}_2\text{S}} = (0.67) \times (0.17) = 0.1139$$


Sol 16: (B)

$$\therefore K_c = \frac{[\text{C}][\text{D}]}{[\text{A}][\text{B}]}$$

$$= \frac{2a \times 2a}{a \times a} = 4.$$

Sol 17: (B)

$$K_c = \frac{[\text{NO}_2]^2}{[\text{N}_2\text{O}_4]} = \frac{4 \times (0.05)^2}{0.05} = 4 \times 0.05 = 0.2$$

Sol 18: (A)


$$2x = 10, \quad \therefore x = \frac{10}{2} = 5$$

$$\text{N}_2 = 30 - 5 = 25 \text{ L}$$


$$\text{H}_2 = 30 - 3 \times 5 = 15 \text{ L}$$

$$\text{NH}_3 = 2 \times 5 = 10 \text{ L}$$

$$\text{Sol 19: (B)} \quad \frac{22}{100} \times 3.2 = 0.704$$

Sol 20: (A)

According to Eq. (1) 2 moles of ammonia are present and to produce 2 moles of NH_3 , we need 1 mole of N_2 and 3 moles of H_2 , hence, $2 - 1 = 1$ mole of N_2 and $4 - 3 = 1$ mole of H_2 are present at equilibrium in vessel.



$$\text{For } 1 \text{ dm}^3, \quad R = k[\text{SO}_2]^2[\text{O}_2]$$

$$R = k \left[\frac{1}{1} \right]^2 \left[\frac{1}{1} \right] = 1$$

$$\text{For } 2 \text{ dm}^3, \quad R = k \left[\frac{1}{2} \right]^2 \left[\frac{1}{2} \right] = \frac{1}{8}$$

So, the ratio is 8 : 1.

Sol 22: (D) When NaNO_3 is heated in a closed vessel, O_2 is liberated and NaNO_2 is left behind. At equilibrium increasing temperature favours forward reaction

Sol 23: (A) 5 moles of SO_2 and 5 moles of O_2 are allowed to react to form SO_3 in a closed vessel. At the equilibrium stage 60% of SO_2 is used up. The total number of moles of SO_2 , O_2 and SO_3 in the vessel now is 8.5.

Sol 24: (B)

K_{c_1} for $\text{H}_2 + \text{I}_2 \rightleftharpoons 2\text{HI}$ is 50

K_{c_2} for $2\text{HI} \rightleftharpoons \text{H}_2 + \text{I}_2$

$$K_{c_1} = \frac{[\text{HI}]^2}{[\text{H}_2][\text{I}_2]}$$

$$K_{c_2} = \frac{[\text{H}_2][\text{I}_2]}{[\text{HI}]^2}$$

On reversing Eq. (i)

$$\frac{1}{K_{c_1}} = \frac{[\text{H}_2][\text{I}_2]}{[\text{HI}]^2}$$

From Eqs. (ii) and (iii)

$$K_{c_2} = \frac{1}{K_{c_1}} = \frac{1}{50} = 0.02$$

Sol 25: (B) $K_p = K_c (\text{RT})^{\Delta n}$

Here, $\Delta n = 2 - 2 = 0$


Sol 26: (C) $K_p = K_c (\text{RT})^{\Delta n}$

Here, $\Delta n = 3 - 2 = 1$

Sol 27: (B) For reaction, $2\text{SO}_3 \rightleftharpoons \text{O}_2 + 2\text{SO}_2$

Here, $\Delta n = 3 - 2 = 1$, i.e., +ve, thus, K_p is more than K_c

$$\left[\because K_p = K_c (\text{RT})^{\Delta n} \right]$$

Sol 28: (A)

$$\Delta n = 2 - 4 = -2$$

$$K_p = K_c (\text{RT})^{\Delta n}$$

$$\therefore K_p = K_c (\text{RT})^{-2}$$

$$\therefore K_c = \frac{K_p}{(\text{RT})^{-2}} = \frac{1.44 \times 10^{-5}}{(0.082 \times 773)^{-2}}$$

Sol 29: (D) Catalyst does not affect equilibrium constant.

Sol 30: (A) $\text{PCl}_5(\text{g}) \rightleftharpoons \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$

$$\begin{matrix} (1-x) & & x & & x \end{matrix}$$


Total number of moles at equilibrium

$$= (1 - x) + x + x$$

$$= 1 + x$$

$$P_{\text{PCl}_3} = \left[\frac{x}{1+x} \right] \times P$$

Sol 31: (A) Equilibrium constant for the reaction:



$$K_c = \frac{1}{4.9 \times 10^{-2}}$$

And for $2\text{SO}_2 + \text{O}_2 \rightleftharpoons 2\text{SO}_3(\text{g})$


$$K_c = \left(\frac{1}{4.9 \times 10^{-2}} \right)^2$$

$$= \frac{10^4}{(4.9)^2} = 416.49$$

Previous Years' Questions

Sol 1: (D) According to Le-chatelier's principle

Sol 2: K_p (equilibrium constant) is independent of pressure and concentration.

Sol 3: (B)

Initial pressure of NH_3 of mole = 15 atm at 27°C

The pressure of 'a' mole of $\text{NH}_3 = p$ atm at 347°C

$$\therefore \frac{15}{300} = \frac{p}{620}$$

$$\therefore p = 31 \text{ atm}$$

At constant volume and at 347°C , mole \propto pressure


$a \propto 31$ (before equilibrium)

$\therefore a + 2x \propto 50$ (after equilibrium)

$$\therefore \frac{a + 2x}{a} = \frac{50}{31}$$


$$x = \frac{19}{62} a$$

$$\begin{aligned} \therefore \% \text{ of } \text{NH}_3 \text{ decomposed} &= \frac{2x}{a} \times 100 \\ &= \frac{2 \times 19a}{62 \times a} \times 100 = 61.29\% \end{aligned}$$



$$\frac{[\text{H}_2]^2[\text{CO}]}{[\text{CH}_3\text{OH}]} = \frac{0.1 \times 0.1 \times 1}{2} = \frac{0.01}{2} = \frac{10 \times 10^{-3}}{2}$$


Sol 5: (A, B, C, D) According to Le-Chatelier's Principle.



Since reaction is endothermic, forward reaction is favoured by increases in temperature.

$K_p = [\text{PO}_2]^{1/2}$. Thus, addition of NaNO_2 or NaNO_3 does not cause any change in K_p .

Sol 7: (C, D, E) According to Le-Chatelier's principle.



Initial stages (C) = 0 or [D] = 0

So, the value of quotient = $\frac{0}{[\text{A}][\text{B}]} = 0$ and increases with time.

Sol 9: (C) According to Le-Chatelier's principle endothermic reaction favours increases in temperature. However exothermic reaction favours decreases in temperature.





Concentration of solids and liquids is taken to be unity.


Sol 11: (E) Increases in pressure favours melting of ice into water because at higher pressure melting point of ice is lowered.

Sol 12: (C) Equation (c) = Equation (a) + Equation (b)

Thus, $K_3 = K_1 \cdot K_2$




$$K_c^1 = \frac{1}{\sqrt{4 \times 10^{-4}}} = 50$$



$$K_p = K_c (\text{RT})^x$$

$x = \Delta n_g = \text{no. of gaseous moles in product} - \text{no. of gaseous in reactant}$

$$= 1 - \left(1 + \frac{1}{2}\right) = 1 - \frac{3}{2} = \frac{-1}{2}$$



$$= 2494.2 + 8.314 \times 300 \ln 4$$

= Positive


$$\Delta G = \text{RT} \ln \frac{Q}{K}$$

Since, ΔG is positive so, $Q > K$ so reaction shifts in reverse direction.

JEE Advanced/Boards

Exercise 1

Sol 1: The system in the initial stage does not contain SO_2, SO_3 will, thus, decompose to form SO_2 and O_2 until equilibrium is reached. The partial pressure of SO_3 will decrease. Let the decrease in partial pressure be $2x$.



At equilibrium $(2x) \quad (2+x) \quad (1-2x)$

Applying law of mass action.

$$K_p = \frac{(1-2x)^2}{(2x)^2(2+x)} \quad (2+x) \rightarrow 2$$

$$900 = \frac{(1-2x)^2}{8x^2}$$

$$\text{Or} \quad \frac{1-2x}{x} = 84.85$$


$$\text{Or} \quad x = 0.0115 \text{ atm}$$

Thus, the partial pressure at equilibrium are:

$$P_{\text{SO}_2} = 2 \times 0.0115 = 0.023 \text{ atm}$$

$$P_{O_2} = 2 + 0.0115 = 2.0115 \text{ atm}$$

$$P_{SO_3} = 1 - 2 \times 0.0115 = 0.977 \text{ atm}$$

Sol 2:

$$\text{At equilibrium} \quad (1-2x) \quad (2x) \quad (x)$$

Applying law of mass action,

$$K_c = \frac{[O_2][CO]^2}{[CO_2]^2} = \frac{x \times (2x)^2}{(1-2x)^2} = 2.96 \times 10^{-92}$$

It can be assumed that $1 - 2x \approx 1.0$ as K_c is very small.

$$\text{So, } 4x^3 = 2.96 \times 10^{-92}$$

$$\text{Or } x = 1.95 \times 10^{-31} \text{ mol L}^{-1}$$


$$[CO] = 2x = 2 \times 1.95 \times 10^{-31}$$

$$= 3.90 \times 10^{-31} \text{ mol L}^{-1}$$

Sol 3: Pressure of NH_3 at $27^\circ C = 15 \text{ atm}$ Pressure of NH_3 at $347^\circ C = P \text{ atm}$

$$\frac{P}{620} = \frac{15}{300}$$

$$P = 31 \text{ atm}$$

Let a moles of ammonia be present. Total pressure at equilibrium = 50 atm 

$$\text{At equilibrium} \quad (a-2x) \quad x \quad 3x$$

$$\text{Total moles } a - 2x + x + 3x = a + 2x$$

$$\frac{\text{Initial number of moles}}{\text{Moles at equilibrium}} = \frac{\text{Initial pressure}}{\text{Equilibrium pressure}}$$

$$\frac{a}{(a+2x)} = \frac{31}{50}$$

$$x = \frac{19}{62} a$$

Amount of ammonia decomposed =

$$2x = 2 \times \frac{19}{62} a = \frac{19}{31} a$$

$$\% \text{ of ammonia decomposed} = \frac{19 \times a}{31 \times a} \times 100 = 61.3$$

Sol 4: $2H_2S(g) \rightleftharpoons 2H_2(g) + S_2(g)$

$$\text{At equilibrium} \quad (0.1-x) \quad x \quad x/2$$

$$\text{Molar conc.} \quad \frac{(0.1-x)}{0.4} \quad \frac{x}{0.4} \quad \frac{x}{0.8}$$

$$K_c = \frac{[H_2]^2[S_2]}{[H_2S]^2} = \frac{\left(\frac{x}{0.4}\right)^2 \left(\frac{x}{0.8}\right)}{\left(\frac{0.1-x}{0.4}\right)^2} = 1.0 \times 10^{-6}$$

$$\text{Or } \frac{x^3}{0.8(0.1-x)^2} = 1.0 \times 10^{-6}$$


as x is very small; $0.1 - x \rightarrow 0.1$

$$\frac{x^3}{0.8 \times (0.1)^2} = 1.0 \times 10^{-6}$$

$$x^3 = 8 \times 10^{-9}$$

$$\text{Or } x = 2 \times 10^{-3}$$

$$\text{So, percent dissociation} = \frac{2 \times 10^{-3}}{0.1} \times 100 = 2.0$$

Sol 5:

$$\text{At equilibrium} \quad (1-x) \quad 2x$$

$$x \text{ (degree of dissociation)} = \frac{D-d}{(n-1)d}$$

$$\text{Given, } d = 38.3, D = \frac{\text{Mol. mass of } N_2O_4}{2} = \frac{92}{2} = 46, n = 2$$

$$\text{So, } x = \frac{46 - 38.3}{38.3} = 0.2$$

$$\text{At equilibrium, amount of } N_2O_4 = 1 - 0.2 = 0.8 \text{ mol}$$


$$\text{And amount of } NO_2 = 2 \times 0.2 = 0.4 \text{ mol}$$

$$\begin{aligned} \text{Mass of the mixture} &= 0.8 \times 92 + 0.4 \times 46 \\ &= 73.6 + 18.4 = 92.0 \text{ g} \end{aligned}$$

Since, 92 gram of the mixture contains = $0.4 \text{ mol } NO_2$

So, 100 gram of the mixture contains

$$= \frac{0.4 \times 100}{92} = 0.43 \text{ mol } NO_2$$

Sol 6:

At equilibrium $(1-x)$ x $x/2$

Total moles at equilibrium = $1-x+x+x/2 = \frac{2+x}{2}$

$$p_{AB_2} = \frac{2(1-x)}{(2+x)}P; p_{AB} = \frac{2x}{(2+x)}P; p_{B_2} = \frac{x}{(2+x)}P;$$

$$K_p = \frac{(p_{AB})^2 (p_{B_2})}{(p_{AB_2})^2} = \frac{\left[\frac{2x}{(2+x)}P\right]^2 \left(\frac{x}{2+x}\right)P}{\left(\frac{2(1-x)}{(2+x)}P\right)^2}$$


$$= \frac{x^3 P}{(2+x)(1-x)^2}$$

As x is very small, $(2+x) \rightarrow 2$ and $(1-x) \rightarrow 1$.

$$\text{So, } K_p = \frac{x^3 P}{2}$$

$$\text{Or } x^3 = \frac{2K_p}{P}$$


$$\text{Or } x = \left(\frac{2K_p}{P}\right)^{1/3}$$

Sol 7:

At equilibrium 0.7 0.3 atm

$$K_p = \frac{(p_{NO_2})^2}{p_{N_2O_4}} = \frac{0.3 \times 0.3}{0.7} = 0.1285 \text{ atm}$$

Let the degree of dissociation of N_2O_4 be x when total pressure is 10 atmosphere.



At equilibrium $(1-x)$ $2x$

Total number of moles = $1-x+2x = 1+x$

$$p_{N_2O_4} = \frac{(1-x)}{(1+x)} \times 10; p_{NO_2} = \frac{2x}{(1+x)} \times 10$$

$$K_p = 0.1285 = \frac{\left(\frac{2x}{1+x}\right)^2 \times 10^2}{\left(\frac{1-x}{1+x}\right) \times 10} = \frac{40x^2}{1-x^2}$$

Since, x is very small, $(1-x^2) \rightarrow 1$

$$\text{So, } x^2 = \frac{0.1285}{40}$$


$$\text{Or } x = 0.0566$$

$$p_{N_2O_4} = \frac{(1-x)}{(1+x)} \times 10 = \frac{1-0.0566}{1+0.0566} \times 10 = \frac{0.9436 \times 10}{1.0566}$$

$$= 8.93 \text{ atm}$$

$$p_{NO_2} = \frac{2x}{(1+x)} \times 10 = \frac{2 \times 0.0566}{1+0.0566} \times 10 = \frac{0.1132}{1.0566} \times 10$$

$$= 1.07 \text{ atm}$$

Sol 8:

At equilibrium $(1-x)$ $(3-3x)$ $2x$

Total number of moles = $1-x+3-3x+2x = 4-2x$

$$p_{N_2} = \frac{(1-x)}{(4-2x)}P; p_{H_2} = \frac{(3-3x)}{(4-2x)}P; p_{NH_3} = \frac{2x}{(4-2x)}P$$

$$K_p = \frac{(p_{NH_3})^2}{p_{N_2} \times (p_{H_2})^3} = \frac{\left(\frac{2x}{4-2x}P\right)^2}{\left(\frac{1-x}{4-2x}P\right) \left(\frac{3-3x}{4-2x}P\right)^3}$$

$$= \frac{4x^2(4-2x)^2}{(1-x) \times 27 \times (1-x)^3 P^2}$$

$$1.6 \times 10^{-5} = \frac{16}{27} \times \frac{x^2(2-x)^2}{(1-x)^4 \times (200)^2}$$

$$\text{Or } \frac{x^2(2-x)^2}{(1-x)^4} = \frac{1.6 \times 10^{-5} \times 27 \times (200)^2}{16}$$

$$= \frac{16 \times 10^{-6} \times 27 \times (200)^2}{16}$$

$$\text{Or } \frac{x(2-x)}{(1-x)^2} = 200 \times 10^{-3} \times \sqrt{27} = 1.039$$

$$\text{Or } x = 0.30$$

Moles of ammonia formed = $2 \times 0.30 = 0.60$

Total moles at equilibrium =

$$(4-2x) = (4-2 \times 0.30) = 3.40$$

$$\% \text{ of NH}_3 \text{ at equilibrium} = \frac{0.60}{3.40} \times 100 = 17.64$$

Sol 9: The volume of SO₂ and O₂ at equilibrium
= (100 - 87) = 13 mL

$$\text{Volume of SO}_2 = \frac{2}{3} \times 13 = 8.67 \text{ mL}$$

$$\text{Volume of oxygen} = \frac{1}{3} \times 13 = 4.33 \text{ mL}$$

$$p_{\text{SO}_3} = \frac{87}{100} \times 1 = 0.87 \text{ atm}$$

$$p_{\text{SO}_2} = \frac{8.67}{100} \times 1 = 0.0867 \text{ atm}$$

$$p_{\text{O}_2} = \frac{4.33}{100} \times 1 = 0.0433 \text{ atm}$$

$$K_p = \frac{p_{\text{SO}_3}}{p_{\text{SO}_2} \times (p_{\text{O}_2})^{1/2}} = \frac{0.87}{(0.0867) \times (0.0433)^{1/2}}$$

$$= \frac{0.87}{(0.0867) \times 0.208} = 48.24 \text{ atm}^{-1/2}$$

Sol 10: Let the molecular mass of the mixture at equilibrium be M_{mix}.

Applying the relation,

$$M_{\text{mix}} = \frac{dRT}{P} = \frac{0.925 \times 0.0821 \times 900}{1} = 68.348$$

Molecular mass of SO₃ = 80

$$\text{Vapour density of SO}_3, D = \frac{80}{2} = 40$$

$$\text{Vapour density of mixture, } d = \frac{68.348}{2} = 34.174$$

Let the degree of dissociation be x.

$$x = \frac{D - d}{(n-1)d} = \frac{40 - 34.174}{\left(\frac{3}{2} - 1\right) \times 34.174} = \frac{5.826 \times 2}{34.174} = 0.34$$

Or x = 34 % dissociated

i.e., SO₃ is 34% dissociated.

Sol 11: K_p = p_{CO₂} = 1

$$\log K_p = 7.282 - \frac{8500}{T}$$

$$\log 1 = 7.282 - \frac{8500}{T}$$

$$T = \frac{8500}{7.282} = 1167.26 \text{ K} = 894.26^\circ\text{C}$$

$$\text{Sol 12: Reaction Quotient} = \frac{[p_{\text{NO}_2}]^2}{p_{\text{N}_2\text{O}_4}} = \frac{100}{10} = 10$$

$$\Delta G^\circ_{\text{reaction}} = 2 \Delta G^\circ_{\text{f NO}_2} - \Delta G^\circ_{\text{f N}_2\text{O}_4}$$

$$= 2 \times 50 - 100 = 0$$


We know that, $\Delta G = \Delta G^\circ - 2.303 RT \log Q_p$

$$= 0 - 2.303 \times 8.314 \times 298 \log 10$$

$$= -5705.8 \text{ J} = -5.705 \text{ kJ}$$

Negative value shows that reaction will be in forward direction.

Sol 13: Ist experiment:



$$K_p = \frac{p_{\text{C}_3\text{H}_5\text{I}} \times p_{\text{HI}}}{p_{\text{I}_2} \times p_{\text{C}_3\text{H}_6}} = \frac{1.8 \times 1.8}{22.1 \times 504} = 2.9 \times 10^{-4}$$

Similarly, solve for second experiment.

$$\text{Sol 14: We know that, } \log \frac{K_2}{K_1} = \frac{\Delta H}{2.303 R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$$

$$\log \frac{0.144}{1.64} = \frac{\Delta H}{2.303 \times 1.987 \times 10^{-3}} \left(\frac{1}{673} - \frac{1}{773} \right)$$

$$\Delta H = -25.14 \text{ kcal for 2 mole}$$

$$= -12.57 \text{ kcal mol}^{-1}$$

Sol 15: $\text{NH}_4\text{HS}(\text{s}) \rightleftharpoons \text{NH}_3(\text{g}) + \text{H}_2\text{S}(\text{g})$

$$\text{Moles of NH}_4\text{HS} = \frac{3.06}{51} = 0.06$$

Degree of dissociation = 0.3

At equilibrium,

$$[\text{NH}_3(\text{g})] = \frac{0.3 \times 0.06}{2}; [\text{H}_2\text{S}(\text{g})] = \frac{0.3}{2} \times 0.06$$

$$K_c = [\text{NH}_3(\text{g})][\text{H}_2\text{S}(\text{g})] = \frac{0.3 \times 0.06 \times 0.3 \times 0.06}{2 \times 2}$$


$$= 8.1 \times 10^{-5}$$

Now applying,

$$K_p = K_c (RT)^{\Delta n} = 8.1 \times 10^{-5} \times (0.082 \times 300)^2 = 0.049$$

Since, NH_4HS is solid, so it causes no change in equilibrium.

Sol 16:



At equilibrium (mole) $(0.1 - x)$ x x

Total number of moles,

$$n = (0.1 - x) + x + x = (0.1 + x)$$

$$PV = nRT$$

$$1 \times 8 = (0.1 + x) \times 0.082 \times 540$$

$$\text{Or } x = 0.08$$

$$K_c = \frac{[\text{PCl}_3][\text{Cl}_2]}{[\text{PCl}_5]} = \frac{x^2}{(0.1 - x) \times 8}$$

$$= \frac{0.08 \times 0.08}{(0.1 - 0.08) \times 8} = 4 \times 10^{-2} \text{ mol L}^{-1}$$

$$K_p = K_c (RT)^{\Delta n} = K_c RT (\Delta n = +1)$$

$$= 4 \times 10^{-2} \times 0.082 \times 540 = 1.77 \text{ atm}$$

Sol 17: We know that,

$$Pm = dRT$$


$$1 \times m = 1.84 \times 0.0821 \times 384$$

$$m = 29 \times 2$$

Vapour density (d) at equilibrium = 29

Initial vapour density = $M / 2 = 92 / 2 = 46$

$$x = \frac{D - d}{(n - 1)d} = \frac{46 - 29}{29} = 0.586$$



$t = 0$	1	0	
t_{eq}	1 - x	2x	(Total moles = 1 + x)

$$P_{\text{N}_2\text{O}_4} = \frac{1 - x}{1 + x} \times P; P_{\text{NO}_2} = \frac{2x}{1 + x} \times P$$

$$K_p = \frac{4x^2P}{1 - x^2} = \frac{4 \times (0.586)^2 \times 1}{1 - (0.586)^2} = 2.09 \text{ atm}$$

Sol 18:



$$2-2a \qquad \qquad 2a \qquad \qquad a$$

$$P = 0.28 \text{ atm}; a = 0.33$$

$$K_p = \frac{\left(\frac{2a}{2+a} \cdot 0.28\right)^2 \left(\frac{a}{2+a} \cdot 0.28\right)}{\left(\frac{2.2a}{2+a} \cdot 0.28\right)^2}$$


$$\text{Sol 19: Partial pressure} = \frac{\text{no. of moles}}{\text{total moles}} \times \text{eq}^m \text{ pressure}$$

Sol 20: Let calculate no. of moles & 0.92 atm is the total equilibrium pressure.

$$k_p = \frac{(P_{\text{H}_2})^2 (P_{\text{CS}_2})}{(P_{\text{CH}_4})(P_{\text{H}_2\text{S}})^2}$$

$$k_p = (\text{no. of moles}) \times \left(\frac{RT}{v}\right)^{\Delta n}$$

Sol 21:




$$\frac{(\text{effusion rate})_{\text{Cl}_2}}{(\text{effusion rate})_{\text{kr}}} = \sqrt{\frac{M_{\text{kr}}}{M_{\text{Cl}_2}}} = \sqrt{\frac{n_{\text{Cl}_2}}{n_{\text{kr}}}}$$

$$\frac{1.16kr}{kr} = \sqrt{\frac{83.1}{71}} = \sqrt{\frac{n_{\text{Cl}_2}}{n_{\text{kr}}}}$$

$$\frac{(1.16)^2 \times n_{\text{kr}}}{83.7} = 71 n_{\text{Cl}_2}; 0.01607 n_{\text{kr}} = 71 n_{\text{Cl}_2}$$

$$n_{\text{kr}} = \frac{71 \times 1}{0.01607} = 4418.17$$



$$\text{Total } P = 1.642 \text{ atm}$$


$$\text{Density} = 1.28 \text{ g/l}$$

$$\alpha = ?$$

$$V.D. = \frac{PRT}{2P} = 1.28x$$

$$\alpha = ?$$

$$V.D. = \frac{PRT}{2P} = 1.28x$$



+

$$\text{eq}^m \quad \frac{3M}{v} + \frac{4M}{v} \rightleftharpoons \frac{x}{v} \quad k_c = \frac{3M}{v}$$

Exercise 2

Single Correct Choice Type

Sol 1: (A) It is an acid-base reaction;

Rest all occurs in reverse direction.

Sol 2: (D) Δn may be zero, positive or negative integers or fractional depending upon nature of reaction.

Sol 3: (A)

$$K_p = K_c (RT)^{\Delta n}$$

$$\therefore \Delta n = -1$$

$$\therefore K_p = 26 \times (0.0821 \times 523)^{-1} = 0.61 \text{ atm}^{-1}$$

Sol 4: (D) $K_p = K_c (RT)^{\Delta n}$; Δn for the given reaction is $2 - 4 = -2$

$$R = 0.082 \text{ L atm K}^{-1} \text{ mol}^{-1}$$

$$\text{Thus, } K_c = \frac{K_p}{(RT)^{\Delta n}} = \frac{1.44 \times 10^{-5}}{(0.082 \times 773)^{-2}}$$

Sol 5: (D) Addition of salt always lowers the vapour pressure. Also vapour pressure of a liquid is its characteristic pressure and depends only on temperature.

Sol 6: (C) An increase in volume for the equilibrium having,

$$K_c = \frac{[\text{SO}_2][\text{Cl}_2]}{[\text{SO}_2\text{Cl}_2]} = \frac{[\text{mole of SO}_2][\text{mole of Cl}_2]}{[\text{mole of SO}_2\text{Cl}_2] \times \text{volume}}$$


Will increase the mole of Cl_2 or SO_2 to maintain K_c values constant.

Sol 7: (C) Only CO_2 is gas at equilibrium and $K_p = P_{\text{CO}_2}$.

Sol 8: (B) Before attaining the equilibrium, the rate of forward reaction decreases due to decrease in concentration of reactants with time and the rate of backward reaction increases due to increase in concentration of products with time.

Sol 9: (C) Le Chatelier's principle is not valid for solid-solid equilibrium.

Sol 10: (B) 1 mole of N_2 reacts with 3 moles of H_2 thus, for



Sol 11: (A) Reversible reactions always attains equilibrium and never go for completion.


Sol 12: (C) Choices (A) and (D) involves precipitation of AgCl and PbI_2 respectively; In (B) H_2 is escaped out.

Sol 13: (D) $\Delta n = (a+b) - (a+b) = 0$

Sol 14: (B) Rest all occur in backward direction.


Sol 15: (A) Le Chatelier proposed a principle to explain the effect of P, T and C on system in equilibrium.

Sol 16: (A)



$$1 - \alpha \quad \alpha \quad \alpha$$

$$\text{For (i) } K_{p_1} = \frac{n_y \cdot n_z}{n_x} \cdot \frac{P_1}{[\sum n]_1} = \frac{\alpha^2}{(1-\alpha)} \cdot \frac{P_1}{(1+\alpha)}$$



$$1 - \alpha \quad 2\alpha$$


$$\text{For (ii) } K_{p_2} = \frac{(n_B)^2}{(n_A)} \times \frac{P_2}{[\sum n]_2} = \frac{4\alpha^2}{(1-\alpha)} \cdot \frac{P_2}{(1+\alpha)}$$

$$\therefore \frac{K_{p_1}}{K_{p_2}} = \frac{\alpha^2 \times P_1}{(1-\alpha) \times (1+\alpha)} \times \frac{(1-\alpha)(1+\alpha)}{4\alpha^2 \times P_2}$$

$$\therefore \frac{P_1}{P_2} = \frac{4 \times 9}{1} = \frac{36}{1}$$

Sol 17: (C) K_c is characteristic constant for a given reaction at a temperature.

Sol 18: (D) K_p remains constant with T. For the reaction,



$$K_c = \frac{4a^2\alpha^2}{(a - a\alpha).V}$$

Where, a is initial mole of N_2O_4 present in V L and α is its degree of dissociation.

$$\text{Also, } K_p = K(RT)^{\Delta n}$$


On reducing the volume of container to $\frac{V}{2}$ L, initial concentration of N_2O_4 becomes $\frac{2a}{V}$. An increase in concentration leads to more dissociation of N_2O_4 in order to have K_c constant, a characteristic constant for a given reaction at a temperature.

Sol 19: (D) K_p is a characteristic constant for a given reaction and changes only with temperature.

Sol 20: (A)

$$K_c = \frac{[\text{NO}_2]^2}{[\text{N}_2\text{O}_4]} = \frac{[1.2 \times 10^{-2}]^2}{4.8 \times 10^{-2}} = 3 \times 10^{-3} \text{ mol L}^{-1}$$

Sol 21: (A)



Given, $2x = 0.2$, Also, $a = 1.1$

$$x = 0.1 \quad b = 2.2$$

$$\begin{aligned} \therefore K_c &= \frac{4x^3}{(a-x)(b-2x)^2} \\ &= \frac{4 \times (0.1)^3}{(1.1-0.1)(2.2-0.2)^2} = 0.001 \end{aligned}$$

Sol 22: (C) K_p is independent of initial concentration.

Sol 23: (A) $\Delta G^\circ = -2.303 RT \log K_p$


$$= -2.303 \times 8.314 \times 300 \log 10^{20}$$

$$= -114.88 \text{ kJ}$$

Sol 24: (A) $K_c = \frac{1}{[\text{O}_2]^5}$; note that the expression does

not involve the concentration terms of solid species.

Sol 25: (D)



$$t = 0 \quad \quad \quad 8 \quad \quad \quad 0 \quad \quad \quad 0$$

$$\text{At equilibrium} \quad (8-a) \quad \quad \quad a/2 \quad \quad \quad \frac{3a}{2}$$


$$\text{Thus, } K_c = \frac{[\text{A}_2][\text{B}_2]^3}{[\text{AB}_3]^2}; \text{ Also, } \frac{a}{2} = 2 \quad \therefore a = 4$$

$$\therefore [\text{AB}_3] = \frac{4}{1}; [\text{A}_2] = \frac{2}{1}; [\text{B}_2] = \frac{6}{1}$$

$$\text{Thus, } K_c = \frac{2 \times 6^3}{4^2} = 27 \text{ mol}^2\text{L}^{-2}$$

Sol 26: (B) K_p and K_c values do not change with catalyst.

Sol 27: (D)




Given, $a - x = 2x$

$$\therefore x = a/3$$

$$\text{Now, } K_c = \frac{[\text{C}]^2[\text{D}]}{[\text{A}][\text{B}]^2}$$

Multiple Correct Choice Type

Sol 28: (A, B, C) According to Le - Chatlier's principle, the yield of product in the reaction



Would be higher at high temperature and low pressure.


Sol 29: (A, B, C, D) Le – Chatlier’s principle.

Sol 30: (C, D) According to Le – Chatlier’s principle, the forward reaction at constant temperature is favoured by introducing an inert gas at constant pressure and by introducing PCl_5 at constant volume.

Comprehension Type

Paragraph 1:

Sol 31: (A)



Pressure increase due to increases in temperature as well as due to increase in moles.

Initially $P \propto T$

$$15 \propto 300$$

$$P \propto 620$$

$$\therefore P = \frac{620}{300} \times 15 = 31 \text{ atm of } 10 \text{ moles of } \text{NH}_3 \text{ at } 620 \text{ K}$$

Now, NH_3 is dissociated to attain 50 atm at 620 K.

Thus, $P \propto n$ or $10 \propto 31$

$$10 + 2x \propto 31$$

$$\therefore 2x = 6.13$$

$$\therefore \alpha = \frac{2x}{10} \times 100 = \frac{6.13 \times 100}{10} = 61.3\%$$

Sol 32: (D)

$$K_p = \frac{n_{\text{N}_2} \times (\text{NH}_2)^3}{(n_{\text{NH}_3})^2} \times \left[\frac{P}{\sum n} \right]^2$$

$$= \frac{\frac{6.13}{2} \times \left[\frac{6.13 \times 3}{2} \right]^3}{[10 - 6.13]^2} \times \left[\frac{50}{10 + 6.13} \right]^2 = 1.528 \times 10^3 \text{ atm}^2$$

Sol 33: (A) $PV = nRT$

$$15 \times V = 10 \times 0.0821 \times 300$$

$$\therefore V = 16.42 \text{ litre}$$

Paragraph 2:

Sol 34: (B) Endothermic reactions are favoured with increase in temperature.

Sol 35: (A) Increase in temperature will favour the change showing absorption of heat.

Sol 36: (C) K_c remains constant and thus ΔG° is not changed.

Previous Years' Questions

Sol 1: (C) When $n_r > n_p$ then $K_p < K_c$

Where n_r = no. of moles of reactant

n_p = no. of moles of product.

Sol 2: (B) The reaction is exothermic so high temperature will favour backward reaction.

Sol 3: (B, C) According to Le-Chatelier's principle.

Sol 4: (A, B, C, D) $\text{C}_2\text{H}_4 + \text{H}_2 \rightleftharpoons \text{C}_2\text{H}_6$ $\Delta H = -32.7 \text{ kcal}$

The above reaction is exothermic, increasing temperature will favor backward reaction, will increase the amount of C_2H_4 . Decreasing pressure will favour reaction in direction containing more molecules (reactant side in the present case). Therefore, decreasing pressure will increase amount of C_2H_4 .

Removing H_2 , which is a reactant, will favour reaction in backward direction, more C_2H_4 will be formed.

Adding C_2H_6 will favour backward reaction and some of the C_2H_6 will be dehydrogenated to C_2H_4 .

Sol 5: (D) $\text{NaNO}_3(\text{s}) \rightleftharpoons \text{NaNO}_2(\text{s}) + \frac{1}{2}\text{O}_2(\text{g})$ $\Delta H > 0$

NaNO_3 and NaNO_2 are in solid state, changing their amount has no effect on equilibrium. Increasing temperature will favour forward reaction due to endothermic nature of reaction. Also, increasing pressure will favour backward reaction in which some $\text{O}_2(\text{g})$ will combine with $\text{NaNO}_2(\text{s})$ forming NaNO_3 .

Sol 6: (A) Temperature and pressure both affects the concentration of X_3Y

Sol 7: (D) $\text{SO}_2\text{Cl}_2(\text{g}) \rightleftharpoons \text{SO}_2(\text{g}) + \text{Cl}_2(\text{g})$

Adding inert gas at constant volume will not affect partial pressure of reactant or products, hence will not affect equilibrium amount of either reactant or products.

Sol 8: (A) Factors affecting equilibrium are pressure, temperature and concentration of product of reactant.

Sol 9: (D) As per Le-Chatelier's principle.

Sol 10: (D) At constant temperature and pressure, chemical reactions are spontaneous in the direction of decreasing Gibbs energy.

Sol 11: (A, B, D) ΔH is dependent on T

(B) K is independent of the initial amount of CaCO_3

(D) ΔH is independent of the catalyst, if any

Sol 12: (B)

	$X_{2(\text{g})} \rightarrow 2X_{(\text{g})}$	
$t = 0$ (No. of moles)	1	0
$t = t$	$1 - \frac{\beta}{2}$	β
$t = t_{\text{eq}}$	$\left(1 - \frac{\beta_{\text{eq}}}{2}\right)$	β_{eq}

$$P_x = 2 \left(\frac{\beta_{\text{eq}}}{1 + \frac{\beta_{\text{eq}}}{2}} \right) \quad n_{\text{Total}} = 1 - \frac{\beta_{\text{eq}}}{2} + \beta_{\text{eq}} = \left(1 + \frac{\beta_{\text{eq}}}{2}\right)$$

$$P_{X_2} = 2 \left(\frac{1 - \beta_{\text{eq}/2}}{1 + \beta_{\text{eq}/2}} \right)$$

$$K_p = \frac{(P_X)^2}{P_{X_2}} = \frac{\left[2 \left(\frac{\beta_{\text{eq}}}{1 + \beta_{\text{eq}}/2} \right) \right]^2}{\left[2 \left(\frac{1 - \beta_{\text{eq}}/2}{1 + \beta_{\text{eq}}/2} \right) \right]^2} = \frac{2\beta_{\text{eq}}^2}{1 - \frac{\beta_{\text{eq}}^2}{4}} = \frac{8\beta_{\text{eq}}^2}{4 - \beta_{\text{eq}}^2}$$

Sol 13: (C) If $\beta_{\text{eq}} = 0.7$

$$K_p = \frac{8 \times (0.7)^2}{4 - (0.7)^2} = \frac{3.92}{3.51} > 1$$

Which can't be possible as $\Delta G^\circ > 0$ ($\Rightarrow K_p < 1$).

\therefore Therefore, option (C) is incorrect.