

The substitution of chlorine atoms into a molecule of alkane results in a compound with anaesthetic properties e.g., chloroform. Increasing the number of chlorine atoms in the compounds increases the depth of anaesthesia given but also increases toxicity. C-F bonds are very stable so their presence leads to non-flammable and unreactive properties. Organofluorine compounds find diverse applications from oil to water repellents to pharmaceuticals, refrigerants and reagents in catalysts.

When C-X carbon is  $sp^3$  hybridised.


**Halogen Derivatives**

When C-X carbon is  $sp^2$  hybridised.

**Allylic**

C-C-C-X

e.g.,



**Alkyl**


$C_nH_{2n+1}X$

e.g.,  $CH_3CH_2CH_2Cl$

**Benzylic**

$C_6H_5CH_2X$

e.g.,



**Vinyl**

C=C-X

e.g.,  $CH_2=CH-Cl$


**Aryl**

Halogen is directly attached to the carbon atom of aromatic ring, e.g.,  $C_6H_5Cl$


**Methods of Preparation**

**(i) Direct halogenation of alkanes:**


Free radical mechanism:



Reactivity order:




**(ii) Addition of HX to alkenes:**





- Unsymmetrical alkenes follow Markovnikov's rule during electrophilic addition.

- If the addition occurs in presence of peroxide, the product will be opposite to Markovnikov's addition (free radical mechanism).


Reactivity order:




**(iii) From alcohols:**



**(iv) Hunsdiecker reaction:**



**(v) Finkelstein reaction:**



**Uses of Some Commercially Important Halogen Derivatives**

**(i) Chloroform ( $CHCl_3$ ):**

- Earlier it was used as anaesthetic but due to its harmful effects it is no longer used for the purpose.
- Used for preparation of chloretone and chloropicrin.
- Used as a solvent for fats, waxes, rubber, resins, etc.

**(ii) Iodoform ( $CHI_3$ ):**

- Used as disinfectant.
- Effective as chemical antiseptic.

**(iii) Freons or chlorofluorocarbons:**

- Used as refrigerants.
- Used as propellant in aerosols such as body spray, hair spray, cleansers, etc.


**(iv) DDT:**

- Used as a powerful insecticide.
- Effective against *Anopheles* mosquitoes which spread malaria.


**(v) Teflon ( $-CF_2-CF_2-$ )<sub>n</sub>:**

- Used as non-stick coating for pans and other cookwares.
- Used in containers and pipework for corrosive chemicals.




**(i) Reduction:**




**(ii) Wurtz reaction:**




**(iii) Reaction with metals:**



**(iv) Corey-House reaction:**



**(v) Oxidation:**



**Chemical Properties**

**Elimination Reactions**

**$S_N1$**

- First order kinetics
- Reactivity:  $3^\circ > 2^\circ > 1^\circ > CH_3X$


**Nucleophilic Substitution Reactions**

**Miscellaneous Reactions**

**$S_N2$**


- Second order kinetics
- Reactivity:  $CH_3X > 1^\circ > 2^\circ > 3^\circ$

**(i) Dehydrohalogenation:**





- Elimination follows the Saytzeff's rule.
- Ease of dehydrohalogenation: Tertiary > Secondary > Primary


**(ii) Action of heat:**



**(I) Hydrolysis with alkalis:**



**(ii) Williamson's synthesis:**



**(iii)  $R-X + KCN \xrightarrow{\text{alc.}} KX + RCN$**

